

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
5 December 2002 (05.12.2002)


PCT

(10) International Publication Number
WO 02/097044 A2

- (51) International Patent Classification⁷: **C12N**
- (21) International Application Number: **PCT/US02/16707**
- (22) International Filing Date: **28 May 2002 (28.05.2002)**
- (25) Filing Language: **English**
- (26) Publication Language: **English**
- (30) Priority Data:
60/293,791 25 May 2001 (25.05.2001) **US**
- (71) Applicant (for all designated States except US):
THOMAS JEFFERSON UNIVERSITY [US/US];
1020 Walnut Street, Philadelphia, PA 19107-5587 (US).
- (72) Inventor; and
- (75) Inventor/Applicant (for US only): **WONG, Albert, J.**
[US/US]; Apartment 3513, 1500 Locust Street, Philadel-
phia, PA 19103 (US).
- (74) Agent: **MONACO, Daniel, A.**; Drinker, Biddle & Reath,
LLP, One Logan Square, 18th and Cherry Streets, Philadel-
phia, PA 19103-6996 (US).
- (81) Designated States (national): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,
MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG,
SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ,
VN, YU, ZA, ZM, ZW.
- (84) Designated States (regional): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR,
GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent
(BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR,
NE, SN, TD, TG).

[Continued on next page]

(54) Title: **ALTERNATIVE SPLICE FORMS OF PROTEINS AS BASIS FOR MULTIPLE THERAPEUTIC MODALITIES**



(57) Abstract: Peptides or antibodies derived from alternative splice forms of proteins associated with a disease or physiologic condition are used as therapeutic or prophylactic agents. Peptides or antibodies derived from alternative splice forms of the vascular endothelial growth factor (VEGF) family of proteins are particularly useful in preventing or delaying the onset of tumors and inducing tumor regression.

WO 02/097044 A2



Published:

— without international search report and to be republished
upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

**ALTERNATIVE SPLICE FORMS OF PROTEINS AS BASIS FOR
MULTIPLE THERAPEUTIC MODALITIES**

Cross Reference to Related Applications

5

This application claims the benefit of co-pending U.S. Provisional Application Serial No. 60/293,791, filed May 25, 2001.

Reference to Government Grant

10

The invention described herein was supported in part by the National Institutes of Health, under grant no. CA69495. The government has certain rights in this invention.

Field of the Invention

15

The present invention relates to the identification and use of alternative splice forms of proteins for the prophylactic or therapeutic treatment of tumors and other disease states.

20

Background of the Invention

Traditionally, vaccines are derived from material completely foreign to the organism being vaccinated. Nevertheless, it is often desirable to immunize
25 an organism with a vaccine based on proteins derived from the organism itself. For example, control of inflammation, prevention of ovulation or other forms of contraception, inhibition of Alzheimer's disease, and prevention or inhibition of tumor growth are all conditions which benefit from immunization with endogenous or "self" proteins.

Peptide vaccines can be used to treat subjects with diseased or abnormal cells; for example, cells infected with viruses, intracellular bacteria or parasites, and tumor cells. The peptide vaccine can induce a cytotoxic T lymphocyte (CTL) response against the diseased or abnormal cells. Cytotoxic T lymphocytes (CTLs) destroy diseased or abnormal cells by direct cytotoxicity, and by providing specific and nonspecific help to other immunocytes such as macrophages, B cells, and other types of T cells. Peptide vaccines can also induce an antibody response, which is useful in the prophylactic and therapeutic treatment of the disease or condition.

Current peptide vaccine technology involves identification of an endogenous normal protein which is associated with the pathogenesis of a given condition. The normal whole protein is then used as the basis for a vaccine. Alternatively, portions of the endogenous protein which are predicted to bind to MHC class I or II motifs are identified and used to produce the vaccine. See Falk et al., *Nature* 351:290, 1991. However, peptide vaccines made only from the partial or whole normal protein sequences can be poorly immunogenic against diseased or abnormal cells, and can also induce an immune reaction against those cells of the body which express the normal protein.

Previous attempts to increase the specific immunogenicity of peptide vaccines have focussed on point mutations in endogenous proteins from various types of cancer cells. These point mutations represent a small area of "non-self" within the larger endogenous protein sequence that may be used to elicit an immune response. However, these point mutations are not effectively recognized by the immune system, and peptide vaccines employing such technology have not produced strong immunologic responses.

Peptide vaccines have also been based on the protein products resulting from gene rearrangements (*i.e.*, deletions, chromosomal rearrangements) that are sometimes present in cancer cells. For example, a chromosome 9:22 translocation in chronic myelogenous leukemia cells produces the BCR/Abl fusion protein. This protein contains an area of "non-self" at the BCR/Abl fusion junction, and has elicited some immunologic response in human patients. However, such gene rearrangements are rare, and chromosomal translocations

are only known to occur in cancer. Thus, the usefulness of vaccines produced from protein products derived from chromosomal rearrangements is limited.

Thus, it is desirable to identify endogenous proteins which are specific to certain tissues in a given disease state, and which also contain immunologic areas of “non-self” that produce a strong immune response, such as a CTL or antibody response. Ideally, such altered endogenous proteins would be commonly occurring and induce little or no cross-reactivity to the corresponding normal protein.

Primary RNA transcripts from certain genes can undergo alternative splicing to produce messenger RNA (mRNA) which differs from the majority of the mRNA produced by the gene. These alternatively spliced mRNAs are translated into alternative splice form proteins that contain different amino acid sequences than the corresponding protein produced by normally spliced mRNA. Alternative splice form proteins are often expressed in a tissue-specific manner under certain physiologic or disease states. Consequently, these alternative splice forms are present in a limited number of cells in a subject suffering from a given disease or condition. For example, it is known that many types of cancer cells produce alternative splice forms which are not found in normal cells from the same subject. Other disease states in which alternative splice forms are specifically produced include diabetes, Alzheimer’s disease and systemic lupus erythematosus (SLE). These alternative splice forms have not heretofore been recognized as a source for vaccines directed against cells from diseased or abnormal tissue which produce the alternative splice form.

Summary of the Invention

It has now been found that peptides derived from alternative splice forms of proteins produced in diseased or abnormal cells are highly immunogenic. Such peptides elicit a strong immune response specific to the “non-self” portion of the peptide. In particular, the peptides can elicit a specific CTL response. The peptides can be used in the therapeutic or prophylactic treatment of a disease or condition which is characterized by the presence of the alternative splice form in certain cells.

The invention thus provides a peptide comprising an amino acid sequence unique to an alternative splice form, wherein the alternative splice form is produced by diseased or abnormal cells of a subject but is substantially absent from the subject's normal cells.

5 The invention also provides a method of identifying immunogenic peptides for treating a subject who has, or is at risk for having, a disease or condition in which diseased or abnormal cells produce at least one alternative splice form, which alternative splice form is substantially absent from normal cells. The method comprises the steps of identifying at least one mRNA which
10 encodes for the at least one alternative splice form; determining at least a partial amino acid sequence of the at least one alternative splice form; and generating at least one peptide comprising an amino acid sequence which is unique to the alternative splice form.

 The invention also provides a method of treating a subject who has, or is
15 at risk for having, a disease or condition in which diseased or abnormal cells produce at least one alternative splice form, which alternative splice form is substantially absent from normal cells. The method comprises administering to the subject an effective amount of at least one peptide comprising an amino acid sequence which is unique to the alternative splice form such that an immune
20 response is generated against the diseased or abnormal cells.

 The invention further provides a method of preventing or delaying the onset of tumor development in a subject at risk for having a tumor in which tumor cells produce at least one alternative splice form, which alternative splice form is substantially absent from non-tumor cells. The method comprises
25 administering to a subject an effective amount of at least one peptide comprising an amino acid sequence which is unique to the alternative splice form, such that an immune response is generated against the tumor cells.

 The invention further provides a method of regressing a tumor in a subject having a tumor in which tumor cells produce at least one alternative
30 splice form, which alternative splice form is substantially absent from non-tumor cells. The method comprises administering to a subject an effective amount of at least one peptide comprising an amino acid sequence which is

unique to the alternative splice form, such that an immune response is generated against the tumor cells.

The invention also provides a method for identifying peptides of the invention which induce MHC-restricted cytotoxic T lymphocyte responses in a
5 subject, comprising obtaining peripheral blood lymphocytes (PBLs), exposing the PBLs to one or more of the peptides such that the PBLs are stimulated, incubating the stimulated PBLs with target cells that either endogenously synthesize the alternative splice form from which the peptide is derived or are pulsed with peptide, and detecting lysis of the target cells.

10 For example, the target cells can be autologously labeled with a radioactive or fluorescent substance. Detecting lysis of the target cells can be accomplished by measuring the amount of autologous label released from the target cells which are lysed by the activated PBLs. Lysis of the target cells can also be detected with an enzyme-linked immunospot ("ELISPOT") assay.

15 The invention also provides antibodies that bind to specific epitopes on alternative splice forms or peptides comprising an amino acid sequence unique to an alternative splice form. The antibodies can be monoclonal or polyclonal, or can be an antibody fragment that is capable of specifically binding to an alternative splice form epitope.

20 The invention also provides a hybridoma that produces a monoclonal antibody which specifically binds alternative splice forms or peptides comprising an amino acid sequence unique to an alternative splice form.

The invention further provides a method of treating a subject having, or at risk for having, a disease or condition in which diseased or abnormal cells
25 produce at least one alternative splice form, which alternative splice form is substantially absent from normal cells. The method comprises administering to a subject an effective amount of at least one antibody specific to an amino acid sequence unique to the alternative splice form, such that one or more clinical symptoms in the subject are ameliorated, or the number of diseased or abnormal
30 cells in the subject is reduced.

Amino Acid Abbreviations

The nomenclature used to describe the peptide compounds of the present invention follows the conventional practice wherein the amino group is presented to the left and the carboxy group to the right of each amino acid residue. In the formulae representing selected specific embodiments of the present invention, the amino-and carboxy-terminal groups, although not specifically shown, will be understood to be in the form they would assume at physiologic pH values, unless otherwise specified. In the amino acid structure formulae, each residue is generally represented by a one-letter or three-letter designation, corresponding to the trivial name of the amino acid, in accordance with the following schedule:

	A	Alanine	Ala
	C	Cysteine	Cys
15	D	Aspartic Acid	Asp
	E	Glutamic Acid	Glu
	F	Phenylalanine	Phe
	G	Glycine	Gly
	H	Histidine	His
20	I	Isoleucine	Ile
	K	Lysine	Lys
	L	Leucine	Leu
	M	Methionine	Met
	N	Asparagine	Asn
25	P	Proline	Pro
	Q	Glutamine	Gln
	R	Arginine	Arg
	S	Serine	Ser
	T	Threonine	Thr
30	V	Valine	Val
	W	Tryptophan	Trp
	Y	Tyrosine	Tyr

Definitions

35


The expression “**amino acid**” as used herein is meant to include both natural and synthetic amino acids, and both D and L amino acids. “**Standard amino acid**” means any of the twenty L-amino acids commonly found in naturally occurring peptides. “**Nonstandard amino acid**” means any amino

-7-

acid, other than the standard amino acids, regardless of whether it is prepared synthetically or derived from a natural source. As used herein, "synthetic amino acid" also encompasses chemically modified amino acids, including but not limited to salts, amino acid derivatives (such as amides), and substitutions.

5 Amino acids contained within the peptides of the present invention, and particularly at the carboxy- or amino-terminus, can be modified by methylation, amidation, acetylation or substitution with other chemical groups which can change the peptide's circulating half life without adversely affecting their biological activity. Additionally, a disulfide linkage may be present or absent in
10 the peptides of the invention.

Amino acids have the following general structure:



15

Amino acids are classified into seven groups on the basis of the side chain R: (1) aliphatic side chains, (2) side chains containing a hydroxylic (OH) group, (3) side chains containing sulfur atoms, (4) side chains containing an acidic or amide group, (5) side chains containing a basic group, (6) side chains
20 containing an aromatic ring, and (7) proline, an imino acid in which the side chain is fused to the amino group.

"Antibody" as used herein includes polyclonal and monoclonal antibodies, chimeric, single chain, chimeric and humanized antibodies, as well as Fab fragments, including the products of an Fab or other immunoglobulin
25 expression library.

The term "**humanized antibody**" refers to an antibody that has its complementary determining regions (CDRs) derived from a non-human species immunoglobulin, and the remainder of the antibody molecule derived from a human immunoglobulin.

30 The term "**chimeric antibody**" means an antibody comprising a variable region and a constant region derived from different species.

The term “chimeric humanized antibody” is meant a chimeric antibody in which at least the constant region is human-derived.

“Peptide” and “protein” are used interchangeably, and refer to a compound comprised of at least two amino acid residues covalently linked by peptide bonds or modified peptide bonds (*e.g.*, peptide isosteres). No limitation is placed on the maximum number of amino acids which may comprise a protein or peptide. The amino acids comprising the peptides or proteins described herein and in the appended claims are understood to be either D or L amino acids with L amino acids being preferred. The amino acid comprising the peptides or proteins described herein may also be modified either by natural processes, such as posttranslational processing, or by chemical modification techniques which are well known in the art. Modifications can occur anywhere in a peptide, including the peptide backbone, the amino acid side-chains and the amino or carboxyl termini. It is understood that the same type of modification may be present in the same or varying degrees at several sites in a given peptide. Also, a given peptide may contain many types of modifications. Modifications include acetylation, acylation, ADP-ribosylation, amidation, covalent attachment of flavin, covalent attachment of a heme moiety, covalent attachment of a nucleotide or nucleotide derivative, covalent attachment of a lipid or lipid derivative, covalent attachment of phosphatidylinositol, cross-linking, cyclization, disulfide bond formation, demethylation, formation of covalent cross-links, formation of cystine, formation of pyroglutamate, formylation, gamma-carboxylation, glycosylation, GPI anchor formation, hydroxylation, iodination, methylation, myristoylation, oxidation, proteolytic processing, phosphorylation, prenylation, racemization, selenoylation, sulfation, transfer-RNA mediated addition of amino acids to proteins such as arginylation, and ubiquitination. See, for instance, Proteins - Structure and Molecular Properties, 2nd Ed., T. E. Creighton, W. H. Freeman and Company, New York, 1993 and Wold F, Posttranslational Protein Modifications: Perspectives and Prospects, pgs. 1-12 in Posttranslational Covalent Modification of Proteins, B. C. Johnson, Ed., Academic Press, New York, 1983; Seifter et al., “Analysis for protein modifications and nonprotein cofactors,” *Meth. Enzymol.* (1990) 182: 626-646

and Rattan et al. (1992), "Protein Synthesis: Posttranslational Modifications and Aging," *Ann NY Acad Sci* 663: 48-62.

As used herein, "**reading frame**" means a specific series of codons in a nucleic acid, for example an mRNA, which produce a given polypeptide when
5 translated.

Brief Description Of The Figures

Figure 1A is a photograph of an agarose gel electrophoresis showing the
10 PCR product representing alternative splice form #1 in VEGFD (SEQ ID NO: 77). Shown is the third round of PCR using primer set 3. The strong band of ~ 300 bp (labeled as "VEGFD#1") represents SEQ ID NO: 77; the band of 1082 bp (labeled as "VEGFD") represents the expected normally spliced mRNA. M: molecular weight in bp; HC2: PCR amplification from HC2 20d2/c cells.
15 **Figure 1B** is a photograph of an agarose gel electrophoresis showing the results of the PCR amplification of NIH-3T3 cells. Shown is the third round of PCR using primer set 3. Only the band corresponding to the expected normally spliced mRNA is present (labeled as "VEGFD"). M: molecular weights in bp; NIH: PCR amplification from NIH-3T3 cells.

20 **Figure 2** is a plot of average tumor volumes in mm^3 vs. time in days for NIH-Swiss mice vaccinated with various immunogenic peptides derived from VEGF family alternative splice forms.

Figures 3A and 3B are survival plots of number of tumors per mouse vs. days after birth and total tumor volume in each mouse vs. days after birth,
25 respectively, for MMTV-neu mice treated with immunogenic peptides derived from VEGF family member alternative splice forms. For both figures, nine litter mate female mice either are left untreated ("No treatment"), given GM-CSF only ("GM-CSF"), or given a combination of VEGF family based alternative splice form vaccines with GM-CSF ("Combo"). Each curve on the
30 survival plots represents a single animal.

Figure 4 shows autoradiographs of dot blots measuring the antibody response in MMTV-Neu mice which were A) left untreated ("No Treat"), B) given GMCSF alone ("GM-CSF"), or C) vaccinated with a combination of SEQ

ID NO: 73, SEQ ID NO: 77 and SEQ ID NO: 81 and GM-CSF ("COMBO"). The antibody response in the mice was measured against 1 microgram of BSA, SEQ ID NO: 82 ("EGFRvIII"), SEQ ID NO: 73 ("R-pep"), SEQ ID NO: 81 homodimer ("RC-pep"), SEQ ID NO: 77 ("H-pep"), SEQ ID NO: 79 ("L-pep"), or SEQ ID NO: 75 ("V-pep") spotted onto nitrocellulose membranes.

Figure 5 is a plot showing lytic events identified using the ELISPOT assay with anti-interferon-gamma antibody. Splenocytes isolated from mice inoculated with HC2 20d2/c tumor cells and then immunized with SEQ ID NO: 73 ("R-pep"), SEQ ID NO: 75 ("V-pep") or SEQ ID NO: 82 ("EGFRvIII") plus GM-CSF that showed tumor regression, or from a mouse inoculated with SEQ ID NO: 75 ("V-pep") and GM-CSF that showed no regression were evaluated in the assay. Mice inoculated with GM-CSF only were used as a control ("control").

Detailed Description Of The Invention

Alternative splice forms (also referred to as "alternative splice form proteins") comprise amino acid sequences not found in the corresponding normal proteins. An amino acid sequence found only in the alternative splice form and not in the corresponding normal protein is considered "unique" to the alternative splice form.

Peptides comprising amino acid sequences which are unique to alternative splice forms are therefore highly immunogenic. As used herein, a "normal" protein is any protein produced from mRNA transcripts which comprise the majority of the total mRNA transcripts produced by a given gene. For example, if two mRNA transcripts are produced by a gene, the mRNA transcript which comprises greater than 50% of the total mRNA transcripts produced by the gene is considered a normally spliced mRNA transcript which is translated into the "normal" protein. The mRNA transcript which comprises less than 50% of the total mRNA produced by the gene is considered an alternatively spliced mRNA, and is translated into an alternative splice form. If three or more mRNA transcripts are produced from a gene, the mRNA transcript which is present in the greatest proportion relative to the other mRNA

transcripts is considered a normally spliced mRNA transcript which is translated into the "normal" protein. Those mRNA transcripts which are present in lesser proportions relative the normally spliced mRNA are considered an alternatively spliced mRNA, and are translated into alternative splice forms. By way of
5 example, if a gene produces three mRNA transcripts in the relative proportions of 40:35:25, the mRNA which comprises 40% of the total mRNA transcripts produced is considered the normally spliced mRNA. The two mRNA transcripts comprising, respectively, 35% and 25% of the total mRNA transcripts produced are considered alternative splice forms.

10 For purposes of the invention, it is assumed that the level of normal or alternatively spliced mRNA produced by a gene is directly proportional to the level of protein produced from the mRNA. The amount of alternatively spliced mRNA, and thus of alternative splice form, in a cell is often a relatively small proportion of the total mRNA transcripts produced by the gene. For example,
15 the alternative splice form can represent 10% or less, 5% or less, 1% or less or 0.5% to 0.1% or less of the total output of a given gene.

The relative amount of mRNA transcripts produced from a gene can be measured by techniques well-known in the art; for example by quantitative reverse-transcriptase polymerase chain reaction (RT-PCR) techniques such as
20 those described in Siebert PD (1993), "Quantitative RT-PCR," Clontech Laboratories, Inc., Palo Alto, CA; Carango P et al. (1995), *Ann. Neurol.* 38: 610-617; and Grove DS (1999), *J. Biomolecular Techniques* 10: 11-16, the entire disclosures of which are herein incorporated by reference. Amounts of mRNA transcripts are typically expressed in relative units (*e.g.*, relative
25 fluorescence units), but can be expressed in terms of mass (*e.g.*, micrograms) or moles (*e.g.*, gram molecular weight).

The amount of an alternative splice form in a diseased or abnormal cell (as evaluated either at the protein or mRNA level) can be at least 50% greater than the level of the same alternative splice form which is found in a normal
30 cell. A "diseased" or "abnormal" cell is identified by certain phenotypic abnormalities which are readily recognized by those skilled in the art upon examination of cells or tissue. For example, the pathology and histopathology

of different cancers is described in Cancer: Principles and Practice of Oncology, (3rd edit., DeVita VT, Hellman S, and Rosenberg SA, eds.), 1989, J. B. Lipincott Co., Phila., PA, the entire disclosure of which is herein incorporated by reference.

5 Cells which are tumorigenic or neoplastic can also be identified by certain growth characteristics and morphology exhibited by the cell in culture. Tumorigenic or neoplastic cells are insensitive to contact-induced growth inhibition, and the cells form foci in the culture vessel when cultured for extended periods. Tumorigenic or neoplastic cells also exhibit characteristic
10 morphological changes, disorganized patterns of colony growth, and the acquisition of anchorage-independent growth. Tumorigenic or neoplastic cells also have the ability to form invasive tumors in susceptible animals, which can be assessed by injecting the cells, for example, into athymic mice or newborn animals of the same species using techniques well-known in the art. See, for
15 example, Combes et al. (1999), "*Cell Transformation Assays as Predictors of Human Carcinogenicity: The Report and Recommendations of ECVAM Workshop 39*," *ATLA* 27, 745-767, available at: <http://altweb.jhsph.edu/science/pubs/ECVAM/ecvam39.htm>, the entire disclosure of which is herein incorporated by reference.

20 Histological, cell culture-based, and other techniques for identifying other types of diseased or abnormal cells are also well-known in the art.

 As used herein, an alternative splice form which is "substantially absent from normal cells" means the alternative splice form is not present in the normal cells or is present in a negligible amount, and in any case is not present in more
25 than about 66% of the level found in a diseased or abnormal cell. An immune response directed specifically against diseased or abnormal cells can therefore be generated in a subject by administering one or more peptides comprising amino acid sequences which are unique to the alternative splice form. Immunization with such peptides has the advantage of not eliciting an immune
30 response against normal cells.

 One type of alternative splice form contains a novel amino acid sequence formed by the joining of two normally distant amino acid sequences. This type

of alternative splice form is created when an mRNA is spliced so as to skip all or part of an exonic sequence which is normally translated into the protein, but which leaves the normal reading frame intact. A novel amino acid sequence is thus created at the "splice junction" by the juxtaposition of two "normal" amino acids which were not heretofore adjacent. Amino sequences from the normal protein flank the splice junction. The skipping of exonic sequences may also create a new codon without shifting the reading frame of the mRNA. In this instance, a new amino acid is inserted at the splice junction which is flanked by normal amino acid sequences. Both types of splice junctions are considered immunologic "non-self" sequences.

Another type of alternative splice form contains novel amino acid sequences translated from coding sequences which are not normally present in the mRNA. Such coding sequences are created by alternative splice events in which intronic sequences, or exonic sequences not typically translated into the normal protein, are included in the mRNA. The coding sequences can also be created by alternative splicing events which alter the native reading frame of the mRNA, resulting in translation of a "missense" amino acid sequence downstream of the alternative splice site. The novel amino acid sequences are considered immunologic "non-self" sequences.

Genes which are involved in a particular disease process, and which are likely to produce alternatively spliced mRNAs, often exhibit altered expression patterns in diseased or abnormal cells as compared to cells from normal tissue.

For example, angiogenesis (the formation of new blood vessels) is a critical process for the continued growth of tumors. Some of the most potent factors for angiogenesis identified in tumors are members of the vascular endothelial growth factor (VEGF) protein family, which are upregulated in tumor cells. The VEGF proteins include VEGF, VEGF-B, VEGF-C, VEGF-D, and placental growth factor (PIGF). The mRNA which produces the VEGF proteins can undergo a variety of splicing events.

Many other genes which produce alternatively spliced mRNAs in a disease-specific manner are known in the art, and can be used in the practice of the present invention. Table 1 contains a representative list of such genes, the

diseases or conditions with which they are associated, the alternative splice forms produced by those genes, and peptides of the invention derived from the alternative splice forms.

-15-

Table 1

<u>Disease</u>	<u>Journal reference</u> ¹	<u>Gene affected (category of protein)</u>	<u>Nature of alternative splice and sequence of peptide derived from alternative splice form</u>	<u>GenBank accession number of gene or gene product² and SEQ ID NO. of alternative splice form (full or partial sequence)</u>
Acute promyelocytic leukemia	Proc. Natl. Acad. Sci. (USA) 99:7640-7645 (2002)	Retinoic acid receptor-alpha (signal transduction protein)	Inclusion of 2 nd intron in RAR-alpha gene, in-frame (underlined): NSNHVASGAPVCHNPNLPSWQALGP YGVVVLAPDTWLSRLSSSPGVEGRS CSARETQA [SEQ ID NO: 3]	AC090426
Myeloblastic leukemia	J. Biol. Chem., Vol. 275, Issue 33, 25255-25261	MCL-1 (transcription factor)	Novel sequences after junction: RNHETAFQ^GWVCGVLPQR [SEQ ID NO: 2]	AF203373 [SEQ ID NO: 1]
Response to stress, infection	J. Biol. Chem., Vol. 276, Issue 31, 29037-29044	Interleukin-1 Receptor-associated Kinase (signal transduction)	In-frame deletion of 30 amino acids (residues 514-543). Deleted region is VYERLEKLQAVVAGVPGHLEAASCIPP SRQ [SEQ ID NO 5]	U52112 [SEQ ID NO: 4]

-16-

Thyroid hormone repression	J. Biol. Chem., Vol. 276, Issue 51, 48196-48205	Branched-chain Aminotransferase (metabolism)	36-bp gap (corresponding to nucleotides 1030-1065 of BCATm) located near the carboxyl terminus	U68418 [SEQ ID NO: 6]
Immunity/inflammation	J Biol Chem, Vol. 274, Issue 39, 27857-27866	Human Glucocorticoid Receptor Beta Isoform (signal transduction)	hGRbeta has additional nonhomologous 15 amino acids at C terminus.	
Oncogenic progression	J Biol Chem, Vol. 273, Issue 43, 28208-28218	Bile salt-dependent lipase (BSDL) (body metabolism)	Underlined is novel sequence: APVPPTGDSGAPPVPP [SEQ ID NO: 8]	AF081673 [SEQ ID NO: 7]
Endothelial cell activation	J Biol Chem, Vol. 274, Issue 3, 1415-1422	Variant of <i>HOXA9</i> (transcription factor)	Inframe junction of sequences: DKPPIDP^NNPAA^NW [SEQ ID NO: 10]	U82759 [SEQ ID NO: 9]
Regulation of Folate	J Biol Chem 271:23820-23827	Folylpolyglutamate Synthetase Gene (metabolism)	Alternate exon: AVSARGATTEGPARRGMS [SEQ ID NO: 12]	U33557 [SEQ ID NO: 11]
Cancer	J. Biol Chem, 267: 4732-4739, 1992	CD44 splicing variants (cell adhesion/signal transduction)	Five different splicing variants inserted into CD44 coding sequence, all inframe	M83324 [SEQ ID NOS: 13 & 14] M83325 [SEQ ID NOS: 15 & 16] M83326 [SEQ ID NOS: 17 & 18] M83327 [SEQ ID NOS: 19 & 20] M83328 [SEQ ID NOS: 21 & 22]
Uterine cancer	Cancer Research, Vol. 54, Issue 13 3337-3341	CD44	Containing variant exons v3 to v10, including the v7/v8 transition epitope	

-17-

Thyroid cancer	Cancer Research, Vol. 56, Issue 5 1037-1042	CD44	Contains subsegment from exon 4 joined to a subsegment of exon 13 (v8), followed by the complete sequence of exons 14 and 15	
Gastrointestinal tumors	Cancer Research, Vol. 55, Issue 19 4273-4277	CD44	Retention of intron 9	
Dysplastic, and neoplastic cervical epithelium	Clinical Cancer Research, Vol. 1, Issue 10 1125-1132	CD44 variants		
Melanoma	Clinical Cancer Research, Vol. 2, Issue 3 447-456	CD44 variants		
Atherosclerosis	J Biol Chem 268:17528-17538, 1993	Variant of Human VLDL receptor (lipid metabolism)	Lacks 84 nucleotides	
Allergic response	J Biol Chem 264: 5912-5915	Variants of IgE receptor alpha chain (receptor/signal transduction)	Exon 2 or part of exon 4 missing	

-18-

Cardiac Function	J. Biol. Chem. In press	Sarco/Endoplasmic Reticulum Ca ²⁺ -ATPase (SERCA) 3 isoforms (ion channels)	Three novel splice forms	AF458230 [SEQ ID NO: 23] AF458229 [SEQ ID NO: 24] AF458228 [SEQ ID NO: 25]
Breast Cancer	<i>Cancer Research</i> 59, 2546-2550	<i>BRCA2</i> (DNA repair)	Junction of exon 11 to 13 inframe LILVGEPSI^STPDGTIK [SEQ ID NO: 26]	U43746 [SEQ ID NO: 27]
Prostate Cancer	<i>Cancer Research</i> , Vol. 54, Issue 5 1190-1193	Estrogen receptor	Variant lacking all of exon 4	
Breast Cancer	<i>Cancer Research</i> , Vol. 53, Issue 24 5934-5939	Constitutively active estrogen receptor (hormone metabolism/signal transduction)	Lack of exon 5	
Breast Cancer	<i>Cancer Research</i> , Vol. 56, Issue 19 4324-4327	exon 4-deleted estrogen receptor mRNA, deleted in exons 2-4 or in regions within exons 3-7		
Small Cell lung cancer	<i>Cancer Research</i> 60, 1840-1844	Neuron-restrictive Silencer Factor Repressor (transcriptional regulation)	Novel exon inserted: VGYGYHLVIFTRV [SEQ ID NO: 29]	AF228045 [SEQ ID NO: 28]

-19-

Lung Cancer	Cancer Research, Vol. 57, Issue 11 2256-2267	FHIT gene (nucleotide metabolism)	Various deletions of coding exons or insertions of novel exons	
Breast Cancer	Cancer Research, Vol. 56, Issue 21 4871-4875	FHIT gene (nucleotide metabolism)	Additional 32 amino acids at amino terminus due to splicing	
Breast Cancer	Cancer Research 59, 4190-4193	Steroid Receptor RNA Activator (hormone metabolism)	Deletion of 203 bp between positions 155 and 357 [SEQ ID NO: 31]	AF092038 [SEQ ID NO: 30]
Acute lymphoblastic leukemia	Cancer Research, Vol. 56, Issue 9 2171-2177	ALL-1 gene (transcription factor)	Deletion of exon 8	
Endometrial hyperplasia	Cancer Research, Vol. 58, Issue 12 2500-2503	PTEN gene (signal transduction)	4 bp deletion in exon 8	
Endometrial Cancer	Cancer Research, Vol. 57, Issue 24 5579-5583	Sex hormone binding globulin (hormone metabolism)	Deletion of entire exon VII	
Teratocarcinoma	Cancer Research, Vol. 54, Issue 1 220-225	PDGF-alpha receptor (receptor/signal transduction)	Lacks exon 14	
Colon Cancer	Cancer Research, Vol. 56, Issue 8 1731-1736	DT-diaphorase (drug detoxification)	Deletion of exon 4	
Breast Cancer	Clinical Cancer Research Vol. 6, 1135-1139		p66 isoform	

-20-

Desmoplastic round cell tumors	Clinical Cancer Research Vol. 6, 3522-3529	EWS gene (transcription factor)	Deletion of exons 5, 6, 7 or 8	
Muscle development	Proc. Natl Acad Sci 92:2686-2690	Nicotinic acetylcholine receptor (neurotransmission/signal transduction)	Inframe deletion of exon 5	
Epithelial neoplasias	Cell Growth & Differentiation, Vol. 6, Issue 9 1185-1191	Tuberous sclerosis gene (signal transduction)	Deletion from amino acids 947 to 990, (deletion of 129-bp exon), inclusion of separate 69-bp exon encoding a novel serine-rich amino acid sequence (1272 to 1295)	
Gastric cancer	Molecular and Cellular Biology, Oct. 1996, p. 5518-5526	Ron (tyrosine kinase receptor/signal transduction)	Joining of 2677 to 2825 (inframe deletion of 147 bp exon)	
Breast and ovarian cancer	Molecular and Cellular Biology, 17:444-452	BRCA1 (DNA repair)	Inframe deletion of exon 11	
Phenylketonuria	J. Biol. Chem 266: 9351- 9354, 1991	Phenylalanine Hydroxylase gene (metabolism)	Deletion of Ile at position 94 or 95	
X-linked agammaglobulinemia	J. Biol. Chem. 271:30303-30306	Bruton's tyrosine kinase (signal transduction)	Deletion of 33 amino acids (amino acid residues 48-80)	

-21-

(immunodeficiency disorder)					J05500 [SEQ ID NO: 37]
Hereditary elliptocytosis (anemia)	J. Biol. Chem. 266:15154-15159,1991	Deletion of exon X, out of frame after junction			
Osteogenesis imperfecta	J. Biol. Chem. 266, No. 33, Issue of November 25, pp. 22370-22374	Pro- α 1(I) collagen (connective tissue)	Inframe 9 bp (3 amino acid) deletion GPPGA [^] PGAPG [SEQ ID NO: 38]		
Defective apoptosis	J. Biol. Chem. 273:30139-30146	Bcl-2-related ovarian killer (apoptosis regulation)	Deletion of amino acid 76-118 TVLLRLG [^] ITWGKVV [SEQ ID NO: 39]		
Alzheimer's disease	J. Biol. Chem. 267:10804-10809	Beta-A4 amyloid precursor protein (amyloid protein)	Inframe deletion of exon 15		
Alcoholism	J. Biol. Chem. 276: 29764-29771	N-Methyl-D-aspartate R1 (neurotransmission/signal transduction)	Eight isoforms generated by alternative splicing of exons 5, 21, and 22		

-22-

Goodpasture disease (autoimmunity)	J. Biol. Chem. 275: 40392-40399	Goodpasture antigen-binding protein (serine/threonine kinase, signal transduction)	Deletion from 1519-1596 (amino acids 371-396)	AF232930 [SEQ ID NO: 32] (normal) and [SEQ ID NO: 33] (alternative splice form) AF232935 [SEQ ID NO 34] (alternative splice form partial sequence)
Induction of apoptosis	J Biol Chem. 273:11930-11936	BAX (apoptosis regulation)	Insertion of 49 bp, novel C-terminus: GLPLAESLKRRLMSLSPGRPPLLWDAH VADRDHLCGGSAHRLTHHLEEDGLRP PAALDCVFPP [SEQ ID NO: 35]	
Systemic Lupus Erythematosus	Biochemical and Biophysical Research Communications 272, 877-881	Caspase-8 (apoptosis regulation)	Alternative splicing of intron 8, frameshift after junction (underlined) HLDAGTVEPKREK [SEQ ID NO: 36]	
Diabetes (non insulin dependent diabetes)	Biochemical and Biophysical Research Communications 181:1419-1424	Insulin receptor (receptor/signal transduction)	Alternative splicing of exon 11 resulting in 12 additional amino acids at C-terminus	

-23-

Pelizaeus-Merzbacher Disease	Annals of Neurology 1995, 38, pp. 610-617	DM20 – Myelin Component	Two alternative splice forms of DM20 – Alt 1 is a fragment of 224 bp encompassing nucleotides –20 to +369 in which 162 bp of exon II (from +4 to +166) has been removed, leaving the reading frame intact but substituting Asp for Gly ² . Alt 2 is a fragment of 253 bp from –20 to +369 in which 133 bp (from +33 to +166) is spliced out from exon II, shifting the mRNA out of frame after the splice.	[SEQ ID NOS: 40 & 41] (nucleotide sequence of splice junctions)
-------------------------------------	--	-------------------------	--	---

¹The entire disclosure of each journal article is herein incorporated by reference.²The entire disclosure of each GenBank record is herein incorporated by reference.

As can be seen from Table 1, certain types of diseases or conditions tend to produce diseased or abnormal cells containing alternative splice forms. For example, alternative splice forms are present in diseased or abnormal cells caused by infections or stress; cancers (*e.g.*, acute promyelocytic leukemia; acute lymphoblastic leukemia; myeloblastic leukemia; uterine cancer; thyroid cancer; gastrointestinal tumors; dysplastic and neoplastic cervical epithelium; melanoma; breast cancer; prostate cancer; lung cancer; endometrial cancer; teratocarcinoma; colon cancer; desmoplastic round cell tumors; epithelial neoplasias; gastric cancer; ovarian cancer); disorders or conditions of the immune system (*e.g.*, allergic response, x-linked agammaglobulinemia, immunity/inflammation, systemic lupus erythematosus, Goodpasture disease); metabolic disorders (*e.g.*, phenylketonuria, non-insulin dependent diabetes); collagen disorders (*e.g.*, osteogenesis imperfecta); disorders of the arteries (atherosclerosis); inherited red cell membrane disorders (*e.g.*, hereditary elliptocytosis); thyroid hormone repression; endometrial hyperplasia; Alzheimer's disease; and alcoholism.

Within a particular type of disease or condition, alternative splice forms tend to be produced from certain genes. For example, in cancer, the CD44 gene; steroid hormone receptors genes (such as the estrogen receptor gene) and the FHIT gene produce a variety of alternative splice forms from which peptides of the invention can be derived.

One of ordinary skill in the art can identify other genes which potentially produce alternative splice forms using well-known techniques, including linkage analysis, gene expression array analysis, homology searches, and point mutation analysis, and commercially available computer software which can be used to predict exon usage in a given nucleic acid sequence.

The mRNA transcribed from genes which are involved in a particular disease process can be analyzed for the presence of alternative splice patterns using techniques well-known in the art. Such techniques include reverse transcription-polymerase chain reactions (RT-PCR), northern blotting and in-situ hybridization. Techniques for analyzing mRNA sequences are described, for example, in Busting SA (2000), *J. Mol. Endocrinol.* 25: 169-193, the entire

-25-

disclosure of which is herein incorporated by reference. Representative techniques for identifying alternatively spliced mRNAs are also described below.

For example, databases that contain nucleotide sequences related to a given disease gene can be used to identify alternatively spliced mRNA. Databases which include nucleotide sequences include GenBank, Embase, and the Cancer Genome Anatomy Project (CGAP) database. The CGAP database, for example, contains expressed sequence tags (ESTs) from various types of human cancers. An mRNA or gene sequence can be used to query such a database to determine whether ESTs representing alternatively spliced mRNAs have been found for a particular gene expressed in a given disease state.

A technique called "RNase protection" can also be used to identify alternatively spliced mRNAs. RNase protection involves translation of a gene sequence into synthetic RNA, which is hybridized to RNA derived from diseased or abnormal cells. The hybridized RNA is then incubated with enzymes that recognize RNA:RNA hybrid mismatches. Smaller than expected fragments indicate the presence of alternatively spliced mRNAs. The putative alternatively spliced mRNAs can be cloned and sequenced by methods well known to those skilled in the art.

The technique of reverse transcription coupled with the polymerase chain reaction ("RT-PCR") can also be used to identify alternatively spliced mRNAs. In RT-PCR, mRNA from the diseased tissue is converted into cDNA by the enzyme reverse transcriptase, using methods well-known to those of ordinary skill in the art. The entire coding sequence of the cDNA is then amplified via PCR using a forward primer located in the 3' untranslated region and a reverse primer located in the 5' untranslated region. The amplified products can be analyzed for alternative splice forms, for example by comparing the size of the amplified products with the size of the expected product from normally spliced mRNA. A preferred method of determining the relative size of the amplified products is gel electrophoresis through agarose gels. Any change in the size of the amplified product can indicate alternative splicing.

If the first round of PCR yields an indistinct product band, for example when analyzed by agarose gel electrophoresis, a portion of the initial PCR reaction can be used in second round of amplification. The second round of amplification preferably employs a set of primers that are internal to the first set of primers. This process, called "nested PCR," is well known to those of ordinary skill in the art. If the amplified products still produce an indistinct band on an agarose gel after the second round of amplification, a third round of nested PCR may be performed. Once a distinct band representing the amplification product is produced, the band is excised from the gel, and the DNA is extracted and sequenced according to known techniques (*e.g.*, the dideoxy-chain termination method according to Sanger et al. (1977), *Proc. Natl. Acad. Sci. USA* 74: 5463, the entire disclosure of which is herein incorporated by reference).

Once an alternatively spliced mRNA is identified, the amino acid sequence of all or part of the alternative splice form encoded by the mRNA can be determined by translating the mRNA sequence according to known techniques. The amino acid sequence encoded by the alternatively spliced mRNA can also be predicted from standard codon usage tables. See, *e.g.*, Fig. 9.1 on pg. 214 of Lewin B, *Genes VI*, Oxford University Press, Inc., New York, 1997, the entire disclosure of which is herein incorporated by reference. Peptides based on this sequence can then be generated and used in the practice of the invention.

Nucleic acid sequences from a partially sequenced alternatively spliced mRNA (or nucleic acid sequences derived from a partial amino acid sequence of an alternative splice form) can be used to identify peptides of the invention. For example, a probe or primer can be generated from the partial sequence of an alternatively spliced mRNA or alternative splice form. The probe or primer can be used with known molecular biology techniques such as primer extension, nucleic acid sequencing, or nucleic acid hybridization to obtain additional sequence data with respect to the partially sequenced alternatively spliced mRNA. Peptides of the invention can be synthesized from the additional alternatively spliced mRNA sequences obtained in this manner.

-27-

Peptides of the invention must contain at least one amino acid or sequence of amino acids which are not found in the normal protein corresponding to the alternative splice form, or at least one novel juxtaposition of amino acid sequences typically located in different parts of the normal protein. In one embodiment, the peptide of the invention can comprise only amino acid sequences unique to the alternative splice form. Peptides of the invention can be any length, but preferably are between 4 and 50 amino acids in length. Particularly preferred are peptides of the invention which are 7 to 25 amino acids, for example 8 or 9, amino acids in length.

To minimize immunologic cross-reactivity with the corresponding normal protein, it is preferred that the amount of normal amino acid sequence contained in the peptides of the invention is limited. The recognition site for MHC Class I molecules, which are essential for the processing of peptide antigens, is 8 or 9 amino acids in length. Without wishing to be bound by any theory, it is believed that peptides of the invention containing normal amino acid sequences which are 8 or more amino acids in length can generate an unwanted immune response against the normal protein. Therefore, the length of contiguous normal sequence contained within the peptides of the invention is preferably 7 amino acids or less. More preferably, the contiguous normal amino acid sequence is 6, 5, 4, 3, 2 or 1 amino acids in length. It is contemplated that the peptides of the invention can comprise more than one sequence of contiguous normal amino acids.

For peptides of the invention derived from alternative splice forms created by the fusion of ordinarily distant amino acid sequences, it is preferred that the contiguous normal amino acid sequences flanking the splice junction are no more than seven amino acids in length. More preferably, the contiguous normal amino acid sequences flanking the splice junction are each 6, 5, 4, 3, 2 or 1 amino acids in length. In one embodiment, peptides of the invention can comprise contiguous normal amino acid sequences flanking the splice junction of 5 or 6 amino acids in length. Without wishing to be bound by any theory, it is believed that such peptides can produce significantly less cross-reactivity to

the corresponding normal protein, while still having an optimal length for MHC class I antigen presentation.

The peptides of the invention can be prepared by any method for synthesizing peptides. For example, the peptides can be obtained by in vitro translation of the corresponding mRNA. The peptides can also be synthesized in solution or on a solid support in accordance with conventional techniques. Various automatic synthesizers are commercially available and can be used in accordance with known protocols. See, for example, Stewart and Young, *Solid Phase Peptide Synthesis*, 2d. ed., Pierce Chemical Co. (1984); Tam et al., *J. Am. Chem. Soc.* 105:6442, 1983; Merrifield, *Science* 232:341-347, 1986; and Barany and Merrifield, *The Peptides*, Gross & Meienhofer, eds., Academic Press, New York, pp. 1-284 (1979), the entire disclosures of which is incorporated herein by reference.

Alternatively, peptides of the invention can be produced using recombinant DNA technology. For example, nucleic acids encoding the peptides can be synthesized by reverse transcription of the alternatively spliced mRNA, or by chemical synthetic techniques well-known in the art. The nucleic acid coding sequence can then be provided with appropriate linkers and ligated into expression vectors commonly available in the art.

The expression vector can comprise regulatory sequences such as start and stop codons, promoter and terminator regions and an origin of replication. For example, promoter sequences compatible with bacterial hosts can be provided in plasmids containing convenient restriction sites for insertion of the desired coding sequence.

Expression vectors comprising nucleic acids encoding the present immunogenic peptides are preferably transfected into suitable bacterial hosts for expression of the peptides, according to known techniques. Yeast or mammalian cell hosts may also be used, provided the expression vector comprises compatible control sequences. A number of expression vectors and host systems are known in the art, and are commercially available.

Nucleic acids encoding the peptides of the invention can be linked to additional protein coding sequences in an expression vector. Expression of the

linked coding sequences produces a fusion protein comprising the peptide of the invention. The additional coding sequences can comprise sequences which encode other peptides of the invention, or sequences which encode other types of proteins.

5 Other techniques for cloning and expressing nucleic acids are described in, for example, Sambrook et al., *Molecular Cloning, A Laboratory Manual*, Cold Spring Harbor Press, Cold Spring Harbor, N.Y. (1982), and Ausubel et al., (ed.) *Current Protocols in Molecular Biology*, John Wiley and Sons, Inc., New York (1987), and U.S. Pat. Nos. 4,237,224, 4,273,875, 4,431,739, 4,363,877 and
10 4,428,941, the entire disclosures of which are incorporated herein by reference.

 The peptides of the invention can comprise additional amino acids; *i.e.*, amino acids not encoded by the corresponding mRNA. For example, one or more amino acids such as tyrosine, cysteine, lysine, glutamic or aspartic acid can be added to the C- or N-terminus of the peptides. These additional amino
15 acids can be used for: linking two or more peptides of the invention to each other, for coupling one or more peptides of the invention to a polyvalent platform that can enhance presentation of the peptides to the immune system (*e.g.*, polylysine, polyethylene glycol, and the like), for coupling one or more
20 peptides of the invention to a another protein or other molecule; or for modifying the physical or chemical properties of the peptides of the invention. Sites for linking the peptides of the invention to a polyvalent platform or to another protein or other molecule can also be introduced by terminal-NH₂ acylation (*e.g.*, acetylation), thioglycolic acid amidation, terminal-carboxy amidation (*e.g.*, with ammonia or methylamine) of the peptides or biotinylation.

25 Preferably, the peptides of the invention are linked together to form homo- or hetero-multimers. The homo- or hetero-multimers can comprise dimers, trimers, tetramers, pentamers, hexamers or higher multimers. It is understood that formation of the peptides of the invention into multimers should not substantially interfere with ability of the linked peptides to function as
30 desired, *e.g.*, as a cytotoxic T cell determinant or activator of an antibody response.

In a preferred method, homo- or hetero-multimers of the peptides of the invention are formed via cysteine residues which are added to the N- and/or C-terminus of the peptides. The peptides are then formed into multimers via controlled oxidation of the cysteine residues.

5 In another method, a disulfide/amide forming heterobifunctional agent such as N-succidimidyl-3-(2-pyridyl-dithio) proprionate (SPDP) is used to form multimers with the peptides of the invention. For example, the peptides can be linked via formation of a disulfide linkage between a first SPDP functional group and a cysteine residue in one peptide, and formation of an amide linkage
10 between a second SPDP functional group and free amino group in another peptide. Other suitable disulfide/amide forming heterobifunctional agents are known; see, for example, *Immun. Rev.* 62:185, 1982, the entire disclosure of which is herein incorporated by reference.

Multimers can also be formed from the peptides of the invention using
15 bifunctional coupling agents that form a thioether linkage. Suitable thioether forming agents are commercially available, and include reactive esters of 6-maleimidocaproic acid; 2 bromoacetic acid; 2-iodoacetic acid; and 4-(N-maleimido-methyl) cyclohexane-1-carboxylic acid. To effect coupling of the peptides, the carboxyl groups of the thioether forming agents are activated with
20 succinimide or the sodium salt of 1-hydroxy-2-nitro-4-sulfonic acid. A preferred activated thioether coupling agent is succinimidyl 4-(N-maleimidomethyl) cyclohexane-1-carboxylate (SMCC).

It is understood that monomeric or multimeric peptides of the invention can also be linked to other proteins using the techniques and reagents described
25 above.

The peptides of the invention can also be modified as necessary to provide certain desired attributes such as improved pharmacological or immunologic effects, or to facilitate entry of the peptides into cells.

Modifications which can improve immunologic effects of the peptides of
30 the invention include modifications which enhance the CTL or antibody-inducing activity. For example, the hydrophobicity of the peptide N-terminus can be increased, particularly where the second residue of the N-terminal is

already hydrophobic and is implicated in binding to an HLA restriction molecule. Without being bound by any theory, it is believed that increasing the hydrophobicity of the N-terminus of the peptides of the invention enhances the efficiency of presentation to T cells. Thus, the peptides of the invention that
5 contain epitopes for which a host may not generate significant CTL activity can be made CTL-inducing by substituting hydrophobic residues at the N-terminus of the peptide.

Other modifications which improve the immunologic effects of the peptides of the invention include amino acid insertions, deletions, and
10 substitutions (either conservative or non-conservative) which increase the binding affinity of the immunogenic peptide to an MHC molecule for subsequent presentation to a cytotoxic T-lymphocyte. By "conservative substitutions" is meant replacing an amino acid residue with another that is biologically and/or chemically similar; *e.g.*, one hydrophobic residue for
15 another, or one polar residue for another. Combinations of amino acids which are biologically or chemically similar include Gly/Ala; Val/Ile/Leu; Asp/Glu; Asn/Gln; Ser/Thr; Lys/Arg; and Phe/Tyr.

Factors which influence the number and type of amino acid residues that can be substituted or deleted include the spacing between essential epitopic
20 points on the immunogenic peptide, and certain conformational and functional attributes that may be sought (*e.g.*, hydrophobicity vs. hydrophilicity). Replacing one amino acid with another in a given combination would therefore be a conservative substitution. Other conservative substitutions will be apparent to those of ordinary skill in the art.

25 In addition, the contributions made by the side chains of amino acid residues in the peptides of the invention can be probed via systematic replacement of amino acid residues in the peptide with a specified amino acid (*e.g.*, Ala).

The peptides of the invention can also be modified to increase stability
30 of the peptides, either *in vitro* or *in vivo*. Such modifications include synthesizing the peptides to contain at least one D-amino acid. The D-amino acid containing peptides are more resistant to peptidases and are more stable in

serum and tissues compared to their L-peptide counterparts. It is expected that peptides of the invention synthesized as D-amino acid containing peptides will have the same efficacy as the corresponding L-peptide. However, any loss of binding affinity for an MHC molecule would be compensated for by increased
5 *in vivo* stability of the D-amino acid peptide. Stability of an L-amino acid-containing peptide of the invention can also be increased by "capping" the peptide with a D-amino acid, which inhibits exopeptidase destruction of the peptide. D-amino acid containing peptides can also be synthesized as "inverso" or "retro-inverso" forms; that is, by replacing all L-amino acids of a sequence
10 with D-amino acids, or by reversing the sequence of the amino acids and replacing all the L-amino acids with D-amino acids.

Modifications of the peptides of the invention which do not affect the ability of the peptides of the invention to elicit an immune response are also contemplated. For example, amino acid residues which are not required for
15 retention of immunogenic activity can be substituted or deleted. One of ordinary skill in the art can readily determine which amino acids can be substituted or deleted, for example by mutational analysis techniques well-known in the art. Generally, any substitutions, additions or deletions between epitopic and/or conformationally important residues will employ amino acids or
20 moieties chosen to avoid steric and charge interference that might disrupt binding of the peptide to an MHC molecule. Other types of modifications which do not affect the ability of the peptides of the invention to elicit an immune response are discussed *supra* under the definition of "protein" or "peptide."

25 One or more peptides of the invention can be administered to a subject to stimulate an immunologic response against the alternative splice form from which the peptides were derived. The immunologic response stimulated by the peptides includes a cytotoxic T lymphocyte and/or an antibody response. For example, administration of the peptides of the invention can generate an MHC
30 HLA-class I restricted cytotoxic T lymphocyte response, which includes a CD8⁺ T lymphocyte response specific for a target antigen, wherein CD8⁺, MHC class I-restricted T lymphocytes are activated. Administration of the peptides of the

invention can also generate an MHC HLA-class II restricted cytotoxic T lymphocyte response, which includes a CD4⁺ T lymphocyte response specific for a target antigen, wherein CD4⁺, MHC class II-restricted T lymphocytes are activated.

5 As used herein, "subject" includes an animal, preferably a mammal, more preferably a human being, that has or is at risk for having a disease or condition in which diseased or abnormal cells produce the alternative splice form from which immunogenic peptide is derived. The alternative splice form from which the peptide of the invention is derived is substantially absent from
10 normal cells in the subject. Generation of an immunologic response in the host with the peptides results in the prophylactic or therapeutic treatment of the disease or condition associated with the expression of the alternative splice form.

 Preferably, two or more of the peptides of the invention are administered
15 to the subject. For example, two or more peptides can be used which define different epitopes from one or more alternative splice forms. The two or more peptides can be linked to form multimers by the techniques described above, or can be formulated in a composition without forming multimers; *e.g.*, as an admixture. Admixtures of different multimers, or admixtures of multimers and
20 monomeric peptides, are also part of the invention. It is contemplated that admixtures can comprise 1, 2, 3, 4 or 5 or more different monomeric or multimeric peptides of the invention.

 When one type of peptide of the invention is administered as a homomultimer, a plurality of repeating epitopes are presented to the subject's
25 immune system. If two or more different peptides of the invention are administered, either as a heteromultimer or an admixture of peptides, a plurality of heterogeneous epitopes are presented to the subject's immune system. The presentation of a plurality of homo- or heterogeneous epitopes produces a synergistic effect on the subject's immune system. Thus, the immune response
30 which is generated by administering the peptides of the invention as multimers or admixtures is greater than the expected additive effects of each peptide comprising the multimer or admixture.

The peptides of the invention can also be administered to a subject in combination with other peptides that present "T-helper" cell epitopes; *i.e.*, epitopes that stimulate T cells which cooperate in the induction of cytotoxic T cells against the target antigen. Peptides that present "T-helper" cell epitopes are described in Ferrari et al., *J. Clin. Invest.* 88:214-222, 1991, and U.S. Pat. No. 4,882,145, the entire disclosures of which are herein incorporated by reference.

As used herein, a substance which is administered "in combination with" a peptide of the invention can be administered at the same time and in the same site as the peptide (*i.e.*, as a complex or as an admixture), or can be administered at a different time and/or place as the peptide. For example, the substance can be administered in the same site before or after administration of a peptide of the invention, or can be administered simultaneously with a peptide of the invention but in a distant site (*e.g.*, the substance can be administered orally if the peptide is administered parenterally, and vice versa; or the substance can be administered in a contralateral limb if both peptide and the substance are administered parenterally). As used herein, a substance which is "complexed" with a of the invention peptide can be covalently or noncovalently attached to the peptide.

The peptides of the invention can also be administered to a subject in combination with at least one component that primes CTLs. For example, certain lipids are known to prime CTLs *in vivo* against viral antigens. Lipids suitable for priming CTLs include tripalmitoyl-S-glycerylcysteinyl-seryl-serine (P₃CSS), which can effectively prime virus-specific cytotoxic T lymphocytes when covalently attached to a peptide. See Deres et al., *Nature* 342:561-564, 1989, the entire disclosure of which is herein incorporated by reference. In a preferred embodiment, peptides of the invention are complexed to P₃CSS and the resulting lipopeptide is administered to an subject to specifically prime a cytotoxic T lymphocyte response.

The induction of neutralizing antibodies can also be primed with P₃CSS conjugated to a peptide. Thus, peptides of the invention coupled to P₃CSS can also elicit a humoral immune response.

In the practice of the invention, at least one peptide of the invention is administered to a patient in an amount sufficient to elicit an immune response; *i.e.*, an antibody or cytotoxic T lymphocyte response, to the diseased or abnormal cells (the "effective amount"). Where peptides of the invention are administered as a multimer or as an admixture, the effective amount represents the cumulative total of the administered peptides. The presence of an immune response in the subject can be determined, for example, by measuring antigen-specific CTL activity in peripheral blood lymphocytes (PBLs) obtained from the subject during treatment. Alternatively, the presence of an immune response in the subject can be determined by measuring the titer of antibodies specific to the peptides of the invention, or by intradermal injection of the peptides of the invention with subsequent measurement of a delayed-type hypersensitivity (DTH) response.

The effective amount of the peptides of the invention administered to a given subject will depend on factors such as the peptide composition, the manner of administration, the stage and severity of the disease being treated, the weight and general state of health of the subject, and the judgment of the prescribing physician. Generally, an effective amount of the peptides of the invention administered to a 70 kg subject is from about 1 microgram to about 2,000 mg of peptide, preferably about 1 microgram to about 500 mg of peptide, more preferably about 10 micrograms to about 200 mg of peptide, and particularly preferably about 50 micrograms to about 100 mg peptide. An optimal dose is about 500 micrograms of peptide/70 kg subject.

It is understood that the peptides of the invention are often employed in serious disease states; that is, situations where the subject's life is threatened. In such cases, particularly in view of the relatively nontoxic nature of the immunogenic peptides, it is possible to administer substantial excesses of the peptides to the subject; *i.e.*, effective amounts in excess of 2000 mg/70 kg subject.

Single or multiple administrations of an effective amount of the peptides of the invention can be performed with dose levels and patterns selected by the treating physician. For therapeutic use, an initial dose of the peptides is

preferably administered to a subject at the first appearance of clinical symptoms, or shortly after diagnosis of the disease or condition, with "booster doses" administered until at least a partial abatement of symptoms is observed. A preferred dosage regimen for therapeutic treatment comprises administering to a
5 subject an initial dose of from about 10 micrograms to about 100 mg of at least one peptide of the invention, followed by booster dosages every 2-4 weeks of from about 1 microgram to about 1 mg of at least one peptides of the invention for a period of 6 weeks to 3 months, depending on the strength of the subject's immune response or degree of response to the treatment. Other therapeutic
10 dosage regimens are contemplated.

For prophylactic use, an effective amount of at least one peptide of the invention is preferably administered in uniform doses at regular intervals over a period of weeks or months. A preferred dosage regimen for prophylactic use is administration of about 10 micrograms to about 100 mg of at least one peptide
15 of the invention every 2-4 weeks for 6 to 8 months. Other prophylactic dosage regimens are contemplated.

For therapeutic or prophylactic use, the peptides of the invention can be administered by any route which is sufficient to expose the peptides to the subject's immune system. Routes of administration include enteral (*e.g.*, oral,
20 rectal, intranasal, etc.) and parenteral administration. Parenteral administration includes intravenous, intramuscular, intraarterial, intraperitoneal, intravaginal, intravesical (*e.g.*, into the bladder), intradermal, intrapulmonary, inhalation, topical or subcutaneous administration. Also contemplated within the scope of the invention is the instillation of the peptides into the body of the subject in a
25 controlled formulation, with systemic or local release of the peptides to occur at a later time. Preferred routes of administration are via intramuscular, intranasal, intradermal and subcutaneous delivery.

For therapeutic or prophylactic treatment, at least one peptide of the invention can also be administered to a subject via expression of nucleic acid
30 sequences encoding the peptide in cells of the subject. For example, cells of a subject can be infected with an attenuated viral host engineered to express at least one nucleic acid encoding a peptide of the invention. A preferred

attenuated viral host is vaccinia virus. In the practice of the invention, an attenuated viral host comprising nucleic acid sequences encoding one or more of the peptides of the invention is introduced into a subject such that some of the subject's cells are infected with the viral host. The infected cells, under control
5 of the viral host, express the peptides of the invention and elicit an immune response in the subject; *i.e.*, a cytotoxic T lymphocyte or antibody response. Vaccinia viral vectors useful as attenuated viral hosts, and methods for delivering the vectors to a subject are known in the art, for example as described in U.S. Pat. No. 4,722,848, the entire disclosure of which is herein incorporated
10 by reference.

For therapeutic or prophylactic treatment, the peptides of the invention can also be administered to a subject via expression of nucleic acid sequences encoding the peptide in bacterial hosts. In the practice of the invention, a bacterial host comprising one or more nucleic acid sequences encoding a
15 peptide of the invention are introduced into a subject, for example by intradermal or intravesical instillation. The bacterial host expresses the peptides of the invention, and elicits an immune response in the subject; *i.e.*, a cytotoxic T lymphocyte or antibody response. A preferred bacterial host is bacillus Calmette Guerin (BCG), which is described in Stover et al. (Nature 351:456-
20 460, 1991), the entire disclosure of which is herein incorporated by reference. Other bacterial hosts suitable for use in the practice of the invention, such as *Salmonella typhi*, *Listeria monocytogenes* and the like, are known to those skilled in the art.

For therapeutic or prophylactic treatment, the peptides of the invention
25 can also be administered to a subject via expression of nucleic acid sequences encoding the peptide in yeast hosts. In the practice of the invention, a yeast host comprising one or more nucleic acid sequences encoding a peptide of the invention are introduced into a subject, for example orally, or by intradermal or intravesical instillation. The yeast host expresses the peptides of the invention,
30 and elicits an immune response in the subject; *i.e.*, a cytotoxic T lymphocyte or antibody response. Preferred yeast hosts are *Saccharomyces cerevisiae* or *Schizosaccharomyces pombe*.

The peptides of the invention can be administered to a subject in combination with any known carrier or adjuvant. Suitable carriers include keyhole limpet hemocyanin; thyroglobulin; albumins such as human serum albumin; tetanus toxoid; polyamino acids such as poly(D-lysine:D-glutamic acid), and the like. Suitable adjuvants include complete or incomplete Freund's adjuvant, aluminum phosphate, aluminum hydroxide, polylecithins, emulsified oils and alum.

The peptides of the invention can also be administered to a subject in combination with immunostimulatory compounds. For example, the immunogenic peptides can be administered with granulocyte macrophage colony stimulating factor (GM-CSF), which is a cytokine known to enhance the presentation of peptides to dendritic cells for processing and presentation to the immune system. Other suitable immunostimulatory molecules include other cytokines such as IL-12, IL-2, IL-4, IL-5, IL-1alpha, and IL-18; and haptens such as dinitrophenol.

The peptides of the invention can also be administered to a subject by exposing immune system effector cells from a subject to one or more of the peptides *ex vivo*. As used herein, "immune system effector cells" are cells which either prime the immune system to target other cells for elimination, or which effect elimination of targeted cells (*i.e.*, "killer" cells). Immune system effector cells include dendritic cells, lymphokine-activated killer (LAK) cells, natural killer (NK) cells, T-cells and macrophages. Methods of obtaining immune system effector cells, and methods for *ex vivo* treatment of such cells are known in the art, for example as described in Blaese *et al.*, 1995, *Science* 270:475-80; Kohn *et al.*, 1995, *Nature Medicine* 1(10):1017-23; and Ferrari *et al.*, 1992, *Blood* 80:1120-24, the entire disclosures of which are herein incorporated by reference.

For example, immune system cells, or mixtures thereof, can be removed from a subject and maintained in culture. The immune system effector cells can optionally be enriched for a particular cell type, either in culture or upon removal from the subject. The cultured immune system cells are then treated with the immunogenic peptides. Treatment of the cells with peptides of the

invention includes direct exposure of the cells to the peptides, or introduction of a nucleic acid encoding the peptides into the cells. For cells directly exposed to the peptides of the invention, concentrations far in excess of what could be tolerated by a subject if administered *in vivo* can be used. For example, cells
5 can be treated with a 10 micromolar solution of the peptides of the invention, or can be treated with peptides of the invention in a final concentration of about 5 mg/ml to about 10 mg/ml, preferably about 0.1 ng/ml to about 5 mg/ml.

Immune system effector cells are generally capable of internalizing the peptides of the invention upon direct exposure to the peptides in culture.
10 However, the peptides of the invention can be modified so that entry into the immune system effector cells is enhanced. For example, the peptides can be encapsulated in a liposome prior to being administered, as is described in more detail below. The encapsulated peptides are delivered directly into the cells by fusion of the liposome to the cell membrane. Reagents and techniques for
15 encapsulating the peptides of the invention in liposomes are well-known in the art, and include, for example, the ProVectin™ Protein Delivery Reagent from Imgenex.

The peptides of the invention can also be modified to enhance entry into the cells by complexing the peptides with a peptide leader sequence known as a
20 "protein transduction domain" or "PTD." PTDs direct entry of the compound into cells by a process known as "protein transduction." See Schwarze et al. (1999), *Science* 285: 1569 – 1572. PTDs are well-known in the art, and may comprise any of the known PTD sequences including, for example, arginine-rich sequences such as a peptide of nine to eleven arginine residues optionally in
25 combination with one to two lysines or glutamines as described in Guis et al. (1999), *Cancer Res.* 59: 2577-2580, the entire disclosure of which is herein incorporated by reference. Preferred PTDs are sequences of eleven arginine residues or the NH₂-terminal 11-amino acid protein transduction domain from the human immunodeficiency virus TAT protein (SEQ ID NO: 42). Other
30 suitable PTD sequences include other arginine-rich sequences; *e.g.*, 9 or 10 arginines, or six or more arginines in combination with one or more lysines or glutamines. Such leader sequences are known in the art; see, *e.g.*, Guis et al.

(1999), *supra*. A PTD may be located anywhere on the peptides of the invention that does not disrupt the peptides' ability to elicit an immune response, but is preferably located at the C-terminal end.

5 Kits and methods for constructing fusion proteins comprising a peptide of the invention and a PTD are known in the art; for example the TransVector™ system (Q-BIOgene), which employs a 16 amino acid peptide called "Penetratin™" corresponding to the *Drosophila* antennapedia DNA-binding domain; and the Voyager system (Invitrogen Life Technologies), which uses the 38 kDa VP22 protein from Herpes Simplex Virus-1.

10 The peptides of the invention can also be modified to enhance entry into cells by complexing the peptides with heat shock proteins (HSP), for example as described in US Pat. No. 5,935,576 of Srivastava (peptide non-covalently linked to HSP), or Suzue K et al. (1997), *Proc. Natl. Acad. Sci. USA*, 94:13146-13151 (peptide covalently linked to HSP), the entire disclosures of which are herein
15 incorporated by reference. Fusion proteins comprising HSP sequences and a peptide of the invention can also be generated according to standard techniques.

 A nucleic acid encoding a peptide of the invention can be introduced into the immune system effector cells by any known method, for example by transfecting the cells with expression vectors or infecting the cells with an
20 attenuated viral host as described above. Techniques for constructing expression vectors comprising nucleic acid sequences encoding peptides of the invention are discussed above.

 After treatment with at least one peptide of the invention, a portion of the *ex-vivo* treated immune system effector cells can be examined to confirm the
25 presence of appropriate levels of the peptide(s) within the cell, and the remaining treated cells can be reintroduced into the subject. Treated cells may be reintroduced into the subject by parenteral methods, including intravenous infusion and direct injection into the bone marrow. Treated cells are preferably reintroduced into the subject in a saline solution or other pharmaceutically
30 acceptable carrier. The number of treated cells to be reintroduced depends on the purity of the cell population, but a typical dosage is in the range of about 10^5 to about 10^8 cells per kilogram of subject body weight. The number of cells

-41-

available for re-introduction can be increased by expanding the cells in culture prior to treatment with the peptides of the invention.

The invention also provides a method for identifying peptides of the invention which induce MHC-restricted cytotoxic T lymphocyte responses. For example, peripheral blood lymphocytes (PBLs) can be obtained and exposed to one or more peptides of the invention. The ability of the peptides to induce specific cytotoxic activity is determined by incubating the stimulated PBL with autologous labeled (e.g., ^{51}Cr) target cells (such as HLA matched macrophages, T cells, fibroblasts or B lymphoblastoid cells) that endogenously synthesize the targeted antigen (or, alternatively, the cell is pulsed with the peptide of interest), and measuring specific release of label. Preferably, the PBLs are exposed to pools of peptides of the invention, wherein each peptide is about 8 to 20 amino acids long, preferably 9 to 12 amino acids long. Peptides of the invention which induce cytotoxic T lymphocyte activity can be selected from the pools.

Once an peptide of the invention that stimulates a cytotoxic T lymphocyte response is identified, the MHC restriction element of the response can be determined. This involves incubating the stimulated PBL (or short term cultures thereof) with a panel of labeled target cells of known HLA types that have been pulsed with the peptide of interest or the appropriate controls. The HLA allele(s) of cells that are lysed by the stimulated PBL are compared to cells not lysed. The HLA alleles from the lysed target cells represent the MHC restriction element(s) for the cytotoxic T lymphocyte response to the peptide.

The lysis of target cells by stimulated PBLs can be detected either by measuring the release of label from autologously labeled target cells, or with an enzyme-linked immunospot ("ELISPOT") assay. The ELISPOT method is well-known in the art, and kits and reagents for performing the method are commercially available from BD Biosciences Pharmingen, San Jose, CA, 95131-1807. An ELISPOT assay can detect cytokine release from a PBL at the single cell level, allowing the direct determination of cytokine-producing cell frequencies.

Briefly, the ELISPOT method is performed by coating a cell culture dish or well with a cytokine capture antibody, for example an anti-interferon gamma

antibody. The unoccupied spots on the culture dish or well are then blocked with a non-specific binding protein. Target cells, for example tumor cells, that endogenously synthesize the alternative splice form which has been used to stimulate PBLs are added to the culture dish or well. Alternatively, target cells
5 can be pulsed with the peptide of interest. Activated PBLs are then added to the culture dish or well. If the activated PBLs specifically lyse the target cells, the PBLs secrete cytokines such as interferon-gamma, which is captured by the cytokine capture antibody initially coated onto the culture dish or well. A secondary antibody that recognized a different epitope of interferon gamma is
10 then used to detect the presence of any interferon-gamma secreted by the PBLs and captured by the cytokine capture antibody. The secondary antibody can be conjugated with a detectable label, such as fluorescent or radioactive label. Identification of peptides of the invention which induce MHC-restricted cytotoxic T lymphocyte responses by the ELISPOT method is shown in
15 Example 5 below.

The peptides of the invention can be formulated into pharmaceutical compositions, also called vaccine compositions, according to techniques known in the art. Pharmaceutical compositions of the present invention are characterized as being at least sterile and pyrogen-free. As used herein,
20 "pharmaceutical formulations" include formulations for human and veterinary use. The present pharmaceutical formulations may comprise at least one peptide of the invention mixed with a physiologically acceptable carrier medium to form solutions, suspensions or dispersions. Preferred physiologically acceptable carrier media are water, buffered water, normal saline, 0.4% saline, 0.3%
25 glycine, hyaluronic acid and the like.

Pharmaceutical compositions of the invention can also comprise conventional pharmaceutical excipients and/or additives. Suitable pharmaceutical excipients include stabilizers, antioxidants, osmolality adjusting agents, buffers, and pH adjusting agents. Suitable additives include
30 physiologically biocompatible buffers (e.g., tromethamine hydrochloride), additions (e.g., 0.01 to 10 mole percent) of chelants (such as, for example, DTPA or DTPA-bisamide) or calcium chelate complexes (as for example

calcium DTPA, CaNaDTPA-bisamide), or, optionally, additions (e.g., 1 to 50 mole percent) of calcium or sodium salts (for example, calcium chloride, calcium ascorbate, calcium gluconate or calcium lactate). Pharmaceutical compositions of the invention can be packaged for use in liquid form, or can be
5 lyophilized.

For solid compositions, conventional nontoxic solid carriers can be used; for example, pharmaceutical grades of mannitol, lactose, starch, magnesium stearate, sodium saccharin, talcum, cellulose, glucose, sucrose, magnesium carbonate, and the like.

10 Methods for preparing pharmaceutical compounds of the invention are known to those of ordinary skill in the art, for example as described in *Remington's Pharmaceutical Science*, 17th ed., Mack Publishing Company, Easton, Pa. (1985), the entire disclosure of which is herein incorporated by reference.

15 For example, a solid pharmaceutical composition for oral administration can comprise any of the carriers and excipients listed above and 10-95%, preferably 25%-75%, of one or more peptides of the invention. A pharmaceutical composition for aerosol administration can comprise 0.01-20% by weight, preferably 1%-10% by weight, of one or more peptides of the
20 invention in finely divided form along with a surfactant and propellant. Suitable surfactants include esters or partial esters of fatty acids containing from 6 to 22 carbon atoms (e.g., caproic, octanoic, lauric, palmitic, stearic, linoleic, linolenic, olesteric or oleic acid with an aliphatic polyhydric alcohol, or its cyclic anhydride); and mixed esters such as mixed or natural glycerides. The
25 surfactant can comprise 0.1%-20% by weight, preferably 0.25-5% by weight, of the aerosol pharmaceutical composition, with the balance being propellant. A carrier may also be included as desired; e.g., lecithin for intranasal delivery.

The peptides of the invention or pharmaceutical compositions of the invention can be encapsulated in liposomes. Liposomes aid in the delivery of
30 the immunogenic peptides to a particular tissue, such as lymphoid tissue, and can also increase the blood half-life of the peptides or compositions. Liposomes suitable for use in the invention are formed from standard vesicle-forming

lipids, which generally include neutral or negatively charged phospholipids and a sterol, such as cholesterol. The selection of lipids is generally guided by consideration of factors such as the desired liposome size and half-life of the liposomes in the blood stream. A variety of methods are known for preparing liposomes, for example as described in Szoka et al., *Ann. Rev. Biophys. Bioeng.* 9:467, 1980; U.S. Pat. Nos. 4,235,871, 4,501,728, 4,837,028, and 5,019,369, the entire disclosures of which are herein incorporated by reference. Preferably, peptides of the invention are incorporated as part of a liposome, either alone or in conjunction with a ligand molecule that can target the liposome to a particular cell or tissue. Ligands which bind to receptors prevalent in lymphoid cells, such as monoclonal antibodies that bind to the CD45 antigen, are preferred.

The present invention also provides antibodies against alternative splice forms or peptides of the invention. An antibody of the invention specifically binds an epitope of the alternative splice form or peptide of the invention that is not present in the normal protein. The antibody can be a monoclonal antibody, a polyclonal antibody or an antibody fragment that is capable of binding antigen. The antibodies of the invention include chimeric, single chain, and humanized antibodies, as well as Fab fragments and the products of an Fab expression library. Antibody fragments, such as Fab antibody fragments, which retain some ability to selectively bind to the antigen of the antibody from which they are derived, can be made using well known methods in the art. Such methods are generally described in U.S. patent 5,876,997, the entire disclosure of which is incorporated herein by reference.

Polyclonal antibodies of the invention can be produced by immunizing a host with substantially pure alternative splice form or peptide of the invention, using techniques well-known in the art. The antibody titer in the immunized host can be monitored over time by standard techniques, such as with an enzyme linked immunosorbent assay (ELISA) using immobilized peptide. If desired, the antibody molecules can be harvested or isolated from the host (*e.g.*, from the blood or serum) and further purified by known techniques, such as protein A chromatography to obtain the IgG fraction.

At an appropriate time after immunization, *e.g.*, when the specific antibody titers are highest, antibody-producing cells can be obtained from the host and used to prepare monoclonal antibodies by standard techniques, such as the hybridoma techniques described by Kohler and Milstein (1975) *Nature* 5 256:495-497 and by Mishell, B.B. et al., Selected Methods In Cellular Immunology, (Freeman WH, ed.) San Francisco, 1980; the human B cell hybridoma technique (Kozbor et al. (1983) *Immunol. Today* 4:72); the EBV-hybridoma technique (Cole et al. (1985), Monoclonal Antibodies and Cancer Therapy, Alan R. Liss, Inc., pp. 77-96); or trioma techniques. See also Current 10 Protocols in Immunology (1994) Coligan et al. (eds.) John Wiley & Sons, Inc., New York, N.Y.). The disclosures of all citations in this paragraph are herein incorporated by reference in their entirety.

Hybridoma cells producing a monoclonal antibody of the invention can be detected by screening the hybridoma culture supernatants for antibodies that 15 bind the polypeptide of interest, *e.g.*, using a standard ELISA assay. Alternatively, a monoclonal antibody directed against an alternative splice form or peptide of the invention can be identified and isolated by screening a recombinant combinatorial immunoglobulin library (*e.g.*, an antibody phage display library) with the peptide of interest. Kits for generating and screening 20 phage display libraries are commercially available (*e.g.*, the Pharmacia Recombinant Phage Antibody System, Catalog No. 27-9400-01; and the Stratagene SurfZAP Phage Display Kit, Catalog No. 240612). Additionally, methods and reagents suitable for generating and screening an antibody display library can be found in, for example, U.S. Pat. No. 5,223,409; PCT Publication 25 No. WO 92/18619; PCT Publication No. WO 91/17271; PCT Publication No. WO 92/20791; PCT Publication No. WO 92/15679; PCT Publication No. WO 93/01288; PCT Publication No. WO 92/01047; PCT Publication No. WO 92/09690; PCT Publication No. WO 90/02809; Fuchs et al. (1991) *Bio/Technology* 9:1370-1372; Hay et al. (1992) *Hum. Antibod. Hybridomas* 30 3:81-85; Huse et al. (1989) *Science* 246:1275-1281; Griffiths et al. (1993) *EMBO J.* 12:725-734, the entire disclosures of which are herein incorporated by reference.

Recombinant antibodies, such as chimeric and/or humanized monoclonal antibodies, are within the scope of the invention. Such chimeric and humanized (including chimeric humanized) monoclonal antibodies can be produced by recombinant DNA techniques known in the art, for example using methods
5 described in U.S. Pat. No. 5,225,539; PCT Publication No. WO 87/02671; European Patent Application 184,187; European Patent Application 171,496; European Patent Application 173,494; PCT Publication No. WO 86/01533; U.S. Pat. No. 4,816,567; European Patent Application 125,023; Better et al. (1988) *Science* 240:1041-1043; Liu et al. (1987) *Proc. Natl. Acad. Sci. USA* 84:3439-
10 3443; Liu et al. (1987) *J. Immunol.* 139:3521-3526; Sun et al. (1987) *Proc. Natl. Acad. Sci. USA* 84:214-218; Nishimura et al. (1987) *Cancer Res.* 47:999-1005; Wood et al. (1985) *Nature* 314:446-449; Shaw et al. (1988) *J. Natl. Cancer Inst.* 80:1553-1559; Morrison (1985) *Science* 229:1202-1207; Oi et al. (1986) *Bio/Techniques* 4:214; Jones et al. (1986) *Nature* 321:552-525; Verhoeyan et al.
15 (1988) *Science* 239:1534; and Beidler et al. (1988) *J. Immunol.* 141:4053-4060, the entire disclosures of which are herein incorporated by reference.

The present antibodies can be used to isolate peptides of the invention, or the alternative splice forms from which they are derived, using standard techniques such as affinity chromatography or immunoprecipitation. Moreover,
20 the present antibodies can be used to detect alternative splice forms (*e.g.*, in a cellular lysate or cell supernatant) in order to evaluate the abundance and pattern of expression of an alternative splice form.

The present antibodies can also be used diagnostically to monitor protein levels in tissue as part of a clinical testing procedure, for example to determine
25 the efficacy of a given treatment regimen. Detection of protein levels with the present antibodies can be facilitated by coupling the antibody to a detectable substance. Suitable detectable substances include various enzymes (*e.g.*, horseradish peroxidase; alkaline phosphatase; beta-galactosidase; and acetylcholinesterase) prosthetic group complexes (*e.g.*, streptavidin/biotin and
30 avidin/biotin), fluorescent materials (*e.g.*, umbelliferone; fluorescein; fluorescein isothiocyanate; rhodamine; dichlorotriazinylamine fluorescein; dansyl chloride; and phycoerythrin, luminescent materials (*e.g.*, luminol),

bioluminescent materials (*e.g.*, luciferase, luciferin, and aequorin), and radioactive materials (*e.g.*, ^{125}I , ^{131}I , ^{35}S or ^3H).

Antibodies of the invention can also be used in the therapeutic or prophylactic treatment of subjects having a disease or condition in which
5 diseased or abnormal cells produce an alternative splice form having epitopes to which the antibody can bind. The alternative splice form is substantially absent from normal cells in the subject. In the practice of the invention, an effective amount of at least one antibody of the invention is administered to a subject. Preferably, at least one monoclonal antibody specific for an alternative splice
10 form or peptide of the invention are administered to the subject.

An "effective amount" of the present antibodies is the amount which ameliorates one or more clinical symptoms in a subject, or causes a reduction in the number of diseased or abnormal cells in a subject. Amelioration of clinical symptoms would be readily apparent to the ordinarily skilled physician. The
15 number of diseased or abnormal cells in a subject's body can also be readily determined. Suitable techniques for determining the number of diseased or abnormal cells includes direct measurement (*e.g.*, calculating the concentration of leukemic cells in the blood or bone marrow) or by estimation from the size of a tissue mass. As used herein, a "tissue mass" is any localized collection of
20 diseased or abnormal cells in a subject's body, for example a tumor. The size of a tissue mass can be ascertained by direct visual observation or by diagnostic imaging methods such as X-ray, magnetic resonance imaging, ultrasound, and scintigraphy. Diagnostic imaging methods used to ascertain size of a tissue mass can be employed with or without contrast agents, as is known in the art. The
25 size of a tissue mass can also be ascertained by physical means, such as palpation of the tissue mass or measurement of the tissue mass with a measuring instrument such as a caliper. An effective amount of the present antibodies can be, for example, from about 0.1 mg/kg to about 100 mg/kg of body weight, preferably about 50 mg/kg to about 100 mg/kg of body weight, more preferably
30 about 10 mg/kg to about 20 mg/kg of body weight.

-48-

The antibodies of the invention are preferably administered to a subject by parenteral means, for example by intravascular (e.g., intraarterial or intravenous) injection or infusion.

- 5 The following examples are offered by way of illustration, not by way of limitation.

Example 1 - Identification of VEGF Immunogenic peptides

Alternative splice forms of vascular endothelial growth factor (VEGF)
10 proteins were identified in HC2 20d2/c cells, which are NIH-3T3 cells that have been transfected with a constitutively active form of the EGF receptor, EGFRvIII. HC2 20d2/c cells were cultured according to standard techniques. Total RNA from HC2 20d2/c cells was isolated, followed by isolation of poly
15 A+ RNA. The poly A+ RNA was used as a template for first strand cDNA synthesis using oligo dT primers. PCR was then performed with primers from VEGF isoforms. The sequences of the primers used for the PCR are provided below. The primers are represented in the 5' to 3' direction, with nucleotide numbers corresponding to the normal cDNA sequence listed to the left of each primer, where available.

20

Mouse VEGF

Set 1

79-99 CCG AAA CCA TGA ACT TTC TGC (SEQ ID NO: 43)

936-916 CTT GGC GAT TTA GCA GCA GAT (SEQ ID NO: 44)

25

Set 2

117-137 ACC CTG GCT TTA CTG CTG TAC (SEQ ID NO: 45)

909-888 AAA TGG CGA ATC CAG TCC CAC (SEQ ID NO: 46)

-49-

Set 3

126-146 TTA CTG CTG TAC CTC CAC CAT (SEQ ID NO: 47)

815-795 GAA GGA TCT CCT CTT CCT TCA (SEQ ID NO: 48)

5

Murine VEGFB:

Set 1

119-139 CTG CTT GTT GCA CTG CTG CAG (SEQ ID NO: 49)

778-758 TCT GGA AAG CAG CTT GTC ACT (SEQ ID NO: 50)

10

Set 2

155-175 GCC CCT GTG TCC CAG TTT GAT (SEQ ID NO: 51)

739-719 TAC AGG TGA CTG GGT TGA GCT (SEQ ID NO: 52)

Set 3

15

182-202 AGC CAC CAG AAG AAA GTG GTG (SEQ ID NO: 53)

733-713 TGT CTG GGT TGA GCT CTA AGC (SEQ ID NO: 54)

Murine VEGFC:

20

Set 1

151-171 AAC ATG CAC TTG CTG TGC TTC (SEQ ID NO: 55)

1559-1539 CTC TCC CGC AGT AAT CCA CAT (SEQ ID NO: 56)

Set 2

25

292-312 GAG GTC AAG GCT TTT GAA GGC (SEQ ID NO: 57)

1521-1501 CTT GGG CCT CTG TTA CCA TGT (SEQ ID NO: 58)

Set 3

30

301-321 GCT TTT GAA GGC AAA GAC CTG (SEQ ID NO: 59)

1509-1489 TTA CCA TGT GGT CCC ACA GAG (SEQ ID NO: 60)

Murine VEGFD:

	Set 1	
5	15-35 1343-1323	GGA GAA TGC CTT TTG CAA CAC (SEQ ID NO: 61) GCC ATT GCA TGG AAA TGT GGC (SEQ ID NO: 62)
	Set 2	
10	57-77 1234-1214	CAA CTG CTT AGT CAT CGG TAG (SEQ ID NO: 63) ACT TGA CAA AGC AGT GAG CTG (SEQ ID NO: 64)
	Set 3	
15	96-116 1178-1158	ATG TAT GGA GAA TGG GGA ATG (SEQ ID NO: 65) GTT GAA TCA AGG GTT CTC CTG (SEQ ID NO: 66)
	Murine PlGF:	
	Set 1	
		TCT CCT CTG GTA TCA GCG TCT (SEQ ID NO: 67) GCA CTG AAT TCC TGA GTG TCT (SEQ ID NO: 68)
	Set 2	
20	763-743	TGG TGA TTG TGC CTT GAA GGA (SEQ ID NO: 69) TCC ATG CCC CTT ATC ATG GAG (SEQ ID NO: 70)
	Set 3	
25	88-108	TGA AGG ACC TTG GCT CTG GAT (SEQ ID NO: 71) AAT AGA GGG TAG GTA CCA GCA (SEQ ID NO: 72)

PCR was performed with each primer set 1. If a distinct band was visible after this round of PCR, then the band was excised and the PCR product was sequenced using the 5' primer. If the band was not distinct, it was excised, the PCR products was purified, and 10% of the purified PCR product was used for further amplification using primer set 2.

Alternatively, if there were multiple indistinct bands, then the PCR products from the reaction tube were purified using a commercially available kit (Qiagen), and 10% of the purified product was used for another round of PCR using primer set 2. If a band was still indistinct it was excised, the PCR product

-51-

was purified, and 10% of the purified product was is used for further amplification using primer set 3. The distinct bands from the 2nd or 3rd amplification rounds are sequenced using the appropriate 5' primer. Sequences were compared to the known gene sequence to identify alternative splice forms.

5 One alternative splice form was identified in VEGF and VEGFB, and two were found in VEGFD (see Fig. 1 for identification of VEGFD#1; SEQ ID NO: 77 below). Amino acid translation of these sequences provided four VEGF peptides which contain amino acid sequences unique to the VEGF alternative splice forms (hereinafter "VEGF peptides"). The partial coding sequences of
10 the alternatively spliced VEGF mRNAs, and the VEGF peptides encoded by the mRNA partial sequences, are provided below.

VEGF Alt. splice #1

15 R T K P E K C D K P R R (SEQ ID NO:73)
AGA ACA AAG CCA GAA AA^A TGT GAC AAG CCA AGG CGG (SEQ ID NO:74)
505^637

SEQ ID NO: 73 represents the peptide encoded by an alternatively
20 spliced mRNA (SEQ ID NO: 74) spliced to create a junction of nucleotides 505 and 637, bringing together two normally distant sequences. The reading frame of mRNA is preserved, and a new codon is also formed at the splice junction (AAA encoding for a lysine residue).

25

VEGFB Alt. splice #1

V V K Q L V Q T P P L P P (SEQ ID NO:75)
30 GTG GTC AAA CAA CTA GT^G CAG ACG CCG CCG CTT CCT CCA (SEQ ID NO:76)
300^678

SEQ ID NO: 75 represents the peptide encoded by an alternatively
spliced mRNA (SEQ ID NO: 76), spliced to create a junction of nucleotides
300 and 678. The splice junction recreates the native valine at this position, but
35 the subsequent codons are out of frame with the native reading frame of normal VEGFB, so that novel amino acids are produced from nucleotide 679 onward.

VEGFD Alt. splice #1

5 H G P V K M S S F Q E T (SEQ ID NO:77)
 CAT GGA CCA GTG AAG^ATG TCC TCA TTC CAA GAA ACT (SEQ ID NO:78)
 185^752

SEQ ID NO: 77 represents the peptide encoded by an alternatively spliced mRNA (SEQ ID NO: 78), spliced to create a junction of nucleotides 185 and 752. The splice junction shifts the codons out of frame at nucleotide 752 relative to the native reading frame of VEGFD, so that novel amino acids are produced from this point onward.

VEGFD Alt. splice #2

15 L E R S E S C E D R C P (SEQ ID NO:79)
 TTG GAA CGA TCT GAA^AGC TGT GAG GAC AGA TGT CCT (SEQ ID NO:80)
 238^1047

20 SEQ ID NO: 79 represents the peptide encoded by an alternatively spliced mRNA (SEQ ID NO: 80), spliced to create a junction of nucleotides 238 and 1047. The splice junction shifts the codons out of frame at nucleotide 1047 relative to the native reading frame of VEGFD, so that novel amino acids are produced from this point onward.

25 It was confirmed that the VEGF alternative splice forms were specific to the tumor and not found in normal tissue. For the HC2 20d2/c cell model, mRNA from NIH-3T3 cells was subjected to the same PCR conditions using the 3 nested sets of primers described above. NIH 3T3 cells are the cells from which HC2 20d2/c cell line was derived. Either no band or only the normally spliced mRNA was detected for mouse VEGF, VEGFB or VEGFD. Exemplary results are shown in Fig. 1B, which reveals that only the normally spliced VEGFD mRNA was present in NIH-3T3 cells.

30 For the MMTV-neu mouse tumor model (see Example 3, below), normal cells (fibroblasts) were isolated by explant culture as follows. Skin was excised from the mice, trypsinized and then placed into cell culture using DMEM with 5% fetal bovine serum as the culture media. Fibroblasts were allowed to

migrate out from the excised skin and attach to the culture dish, whereupon they were propagated for 5-7 passages. The mRNA from the cultured fibroblasts was harvested, and subjected to the same PCR conditions as the mRNA isolated from MMTV mouse tumor cells using the three nested sets of PCR primers
5 described above. PCR amplification of mRNA from normal MMTV-neu mouse cells revealed no band for the alternative splice form VEGFB, but showed an intact VEGFD band similar to that shown in Fig. 1B, representing the normally spliced mRNA.

The VEGF peptides listed above were synthesized by conventional
10 methods. To avoid potential immune recognition of the normal VEGF family member, the VEGF peptides contained no more than 6 ordinarily contiguous amino acids from the corresponding normal protein. A peptide homodimer was made by adding a cysteine residue to the C-terminus of SEQ ID NO: 73 to give RTKPEKCDKPRRC (SEQ ID NO: 81). Dimerization was accomplished by
15 controlled oxidation of the added C-terminal cysteine residues of SEQ ID NO: 81.

Example 2 - Regression of Tumors by Treatment with VEGF Peptides

20 The VEGF peptides and the SEQ ID NO: 85 homodimer were evaluated for their ability to elicit an anti-tumor immune response in a mouse tumor model. A total of 112 syngeneic mice were divided into eight treatment groups of 14 mice each, and injected with HC2 20d2/c tumor cells. Four days post-injection, the first through fifth treatment groups were vaccinated in the left
25 inguinal area with 100 µg of either SEQ ID NO: 73, SEQ ID NO: 75, SEQ ID NO: 77, SEQ ID NO: 79, or the SEQ ID NO: 81 dimer diluted in 100 µl of PBS mixed with 45 ng of murine GM-CSF as an adjuvant. The total volume of phosphate buffered saline (PBS) was 150 microliters with 2 mg/ml mouse serum albumin as a carrier. The sixth treatment group was vaccinated as above with a
30 combination of all four VEGF peptides and the SEQ ID NO: 81 homodimer (60 µg each) in a total of 200 microliters of PBS with 45 ng of murine GM-CSF and 2 mg/ml mouse serum albumin. The seventh treatment group was vaccinated as

above with 100 µg of a peptide derived from a mutant epidermal growth factor receptor (EGFR) protein called EGFRvIII (SEQ ID NO: 82). The vaccine composition also contained 45 ng of murine GM-CSF and 2 mg/ml mouse serum albumin. The EGFRvIII mutation is the result of a genomic
5 rearrangement in the EGFR gene, and the EGFRvIII peptide (SEQ ID NO: 82) is reported to induce an immune response against tumor cells expressing the mutant receptor. The eighth treatment group was vaccinated with GM-CSF and 2 mg/ml mouse serum albumin in PBS with no peptide, as a control. All treatment groups were then injected once daily for the next 3 days with the 45
10 ng GM-CSF in the original area of vaccination.

Mice that received GM-CSF alone exhibited on average a progression of tumors 5 days after vaccination, and then a moderate level of regression thereafter (Fig. 2, "HC2 only"). This regression is most likely due to spontaneous rejection of the mutant EGFRvIII protein which is expressed in
15 HC2 20d2/c tumor cells. Mice that received the EGFRvIII peptide also exhibited progression of tumors for approximately 5 days, but the progression was not as dramatic as the control mice and there is distinct regression noted thereafter (Fig. 2, "APLEEK"). The regression pattern observed for EGFRvIII vaccination is similar to previous results obtained with this peptide.
20 (Moscattello, D.K., *Cancer Research* 57:1419-1424, 1997).

Fig. 2 shows that all four of the monomeric VEGF peptides and the SEQ ID NO: 81 homodimer induce tumor regression to some degree, which is most notable 12 days after vaccination. For two of the peptides, SEQ ID NO: 75 and SEQ ID NO: 81 (Fig. 2, "VVKQL" and "HGPVK"), regression is seen until ~20
25 days after vaccination; the tumors, however, progress from that point onward. Mice vaccinated with the SEQ ID NO: 73 peptide synthesized without an additional cysteine (Fig. 2, "RTKPEK no C") exhibited a pattern of regression similar to that found for the EGFRvIII peptide. Mice vaccinated with the SEQ ID NO: 81 homodimer or the SEQ ID NO: 79 peptide (Fig. 2, "RTKPEK with
30 C" and "LERSE") show a strong reduction in average tumor volumes by the end of the study, although SEQ ID NO: 81 homodimer treatment group initially shows tumor progression. Mice vaccinated with a combination of all four

-55-

VEGF peptides and SEQ ID NO: 81 homodimer exhibit no tumor progression, and showed the greatest reduction in tumor volume of all treatment groups (Fig. 2, "Combo 5").

These results demonstrate that administering vaccine compositions comprising peptides based on VEGF family alternative splice forms elicit an effective anti-tumor immune response and causes tumor regression. Moreover, administration of a multimer or admixture of the VEGF peptides can further enhance the anti-tumor effect.

10 **Example 3 - Delaying Onset of Breast Cancer Tumors by Treatment With VEGF Peptides**

The ability of the VEGF peptides to prevent or delay the onset of tumors in MMTV-neu mice was evaluated. MMTV-neu mice are transgenic mice carrying an oncogenic form of the human *neu* oncogene under the control of the MMTV promoter, which drives expression of this oncogene in mammary tissue. 100% of female MMTV-neu mice develop multiple mammary tumors within 5-6 months of birth, which progress and eventually require sacrifice of the animal. Analysis of MMTV-neu mouse mammary tumors for VEGF family member alternative splice forms revealed alternative splice forms comprising the VEGF peptides SEQ ID NO: 73 and SEQ ID NO: 77.

Nine littermate MMTV-neu female mice were used in the experiment. One treatment group of three mice were initially vaccinated at 6 weeks of age with a combination of 60 micrograms each of SEQ ID NO: 73, SEQ ID NO: 77, and the SEQ ID NO: 81 homodimer with 45 ng of GM-CSF as the adjuvant. The mice are then vaccinated with the same composition at 5, 6, and 7 months of age. A control group of three mice received 45 ng GM-CSF alone as the adjuvant. A control group of three females received only 45 ng of GM-CSF. Both groups of mice were injected with 45 ng of GM-CSF at the original site of injection for an additional 3 days. This same cycle of vaccination was repeated at 5, 6, and 7 months of age. A second control group of three females received no treatment.

The results are given in Fig. 3, and show that untreated mice or mice given only GM-CSF develop tumors by 158 days after birth. All animals in

-56-

both control groups had tumors by 192 days after birth (Fig. 3A, "GM-CSF" and "No treatment"). In contrast, mice in the treatment group did not develop tumors until 192 days of age, and all animals in the treatment group did not have tumors until 226 days after birth (Fig. 3A, "Combo"). The total tumor volume per animal also revealed a difference in overall tumor burden between the treatment and control groups. By day 240, two of the untreated mice and one of the GM-CSF only mice attained total tumor volumes of greater than 3000 mm³, necessitating sacrifice of the animals (Fig 3B, "GM-CSF" and "No treatment"). However, in the treatment group the greatest tumor burden at day 240 was less than 1200 mm³.

Thus, vaccination with compositions comprising VEGF peptides delayed the onset of tumor formation in a mouse model of human breast cancer, and resulted in a significantly reduced tumor burden.

Example 4 - Antibody response in MMTV-Neu mice vaccinated with a combination of VEGF peptides

A "dot blot" analysis was performed to assess whether an antibody response was generated in MMTV-neu mice which were vaccinated with a combination of VEGF peptides. Three female MMTV mice were injected with a combination of 60 micrograms each of SEQ ID NO: 73, SEQ ID NO: 77, and the SEQ ID NO: 81 homodimer with 45 ng of GM-CSF as the adjuvant (COMBO) as in Example 3. Three female MMTV-neu mice that were left untreated (No Treatment) and three female MMTV-neu mice that were injected with 45 ng GM-CSF alone (GM-CSF) were used as controls. The dot blots were prepared by separately spotting 1 microgram of BSA, SEQ ID NO: 82 ("EGFRvIII"), SEQ ID NO: 73 ("R-pep"), SEQ ID NO: 81 homodimer ("RC-pep"), SEQ ID NO: 81 ("H-pep"), SEQ ID NO: 79 ("L-pep") or SEQ ID NO: 75 ("V-pep") onto nitrocellulose membranes. One membrane was prepared for each experimental and control animal. The membranes were then incubated with serum taken from the appropriate control or experimental animal at 2 months after the last injection, and diluted 1:100. The membranes were then

washed and incubated ^{125}I anti-mouse secondary antibody, washed again and exposed to x-ray film.

The untreated animals (No Treatment; Fig. 4A) or those that received GM-CSF alone (GM-CSF; Fig. 4B) did not show an antibody response against any VEGF or control peptide tested. In contrast, all three animals that had been immunized with SEQ ID NO: 73, SEQ ID NO: 77 and SEQ ID NO: 85 (COMBO; Fig. 4C) showed a strong antibody reaction against SEQ ID NO: 77. In addition, two of the animals demonstrated an antibody response against SEQ ID NO: 73 and SEQ ID NO: 81. However no animal from the immunized group showed an antibody reaction against the BSA control or the three control peptides (EGFRvIII, L-pep, and V-pep) with which the animals were not vaccinated.

Example 5 - Generation of Cytotoxic T lymphocyte Activity by VEGF Peptides

An ELISPOT assay was performed to show that cytotoxic T lymphocyte activity was elicited by VEGF family peptides, as follows. Splenocytes were isolated from mice that were inoculated with HC2 20d2/c tumor cells and subsequently immunized with SEQ ID NO: 73 ("R-pep"), SEQ ID NO: 75 ("V-pep") or SEQ ID NO: 82 ("EGFRvIII") plus GM-CSF that showed tumor regression, or from a mouse inoculated with SEQ ID NO: 75 ("V-pep") and GM-CSF that showed no regression were evaluated in the assay. Splenocytes from mice inoculated with GM-CSF only were used as a control ("control"). HC2 20d2/c tumor cells pulsed with 10 micrograms/ml of the immunizing peptide, or mouse serum albumin (for splenocytes from GM-CSF only treated animals) were used as the target cells for the treatment and control splenocytes, respectively. Untreated HC2 20d2/c tumor cells were used as negative control target cells.

The splenocytes were incubated at various ratios (from 10:1 to 2:1) with target cells in triplicate, and lytic events were identified with an ELISPOT assay using anti-interferon-gamma antibody as the cytokine capture antibody. The specific spots per 10^6 cells were quantitated using the lytic events from

untreated target cells as the background, and then performing linear regression analysis. The results are shown in Fig. 5.

Fig. 5 shows that the mice immunized with either SEQ ID NO: 73 ("R-pep") or SEQ ID NO: 75 ("V-pep") and which had undergone tumor regression had strong CTL responses when target cells were pulsed with the vaccinating peptide. These responses were greater than that seen for the mouse treated with SEQ ID NO: 82 ("EGFRvIII") which had undergone tumor regression. In contrast, splenocytes from the mouse treated with only GM-CSF, or the mouse vaccinated with SEQ ID NO: 73 ("V-pep"), neither of which had undergone tumor regression, showed poor CTL responses.

Example 6 - Generation of Cytotoxic T lymphocyte Activity by Admixtures of VEGF Peptides

An ELISPOT assay is performed to show that cytotoxic T lymphocyte activity is elicited by VEGF family peptides, as follows. Splenocytes are isolated from mice that are inoculated with HC2 20d2/c tumor cells and subsequently immunized with a combination of 60 micrograms each of SEQ ID NO: 73, SEQ ID NO: 77, and the SEQ ID NO: 81 homodimer with 45 ng of GM-CSF as the adjuvant, or SEQ ID NO: 82 ("EGFRvIII") plus 45 ng GM-CSF. Mice inoculated with GM-CSF only are used as a control ("control"). HC2 20d2/c tumor cells pulsed with 10 micrograms/ml of the immunizing peptides, mouse serum albumin (as positive control), or untreated HC2 20d2/c tumor cells (negative control) are used as the target cells. The splenocytes are incubated at various ratios (from 10:1 to 2:1) with target cells in triplicate, and lytic events are identified with an ELISPOT assay using anti-interferon-gamma antibody as the cytokine capture antibody. The specific spots per 10^6 cells are quantitated using the lytic events from untreated target cells as the background, and then performing linear regression analysis.

The experiment is repeated with the following combinations of 60 micrograms each of the VEGF peptides as indicated:

- 1) SEQ ID NO: 73 and SEQ ID NO: 77;
- 2) SEQ ID NO: 77 and SEQ ID NO: 81; and

3) SEQ ID NO: 73; and SEQ ID NO: 81.

Example 7 - Tumor Prevention and Regression with Peptides, Multimeric Peptides, and Peptide Admixtures

5 Peptides of the invention derived from the alternative splice forms associated with breast cancer, ovarian cancer, prostate cancer, lung cancer, skin cancer, lymphoma, bladder cancer, and pancreatic cancer are tested for their ability to either prevent tumors or induce the regression of tumors in experimental mouse models.

10

Mouse Models of Cancer

 The mouse models used in the experiments are those in which the animals endogenously develop the relevant cancers or support the growth and tumorigenic properties of the relevant tumor-derived cell lines. For breast
15 cancer, mice with a heterozygous deficiency for the *pten* gene or MMTV-neu transgenic mice are used; for ovarian cancer, mice with a heterozygous mutation in either BRCA1 or BRCA2 are used; for prostate cancer, transgenic mice with the SV40 early gene under the control of the probasin promoter are used; for lymphoma, mice with a homozygous deficiency for the p53 or p19^{ARF}
20 gene are used; for bladder cancer, mice fed *N*-butyl-*N*-(4-hydroxybutyl)nitrosamine (BBN) are used. Mouse models of lung and bladder cancer can be obtained, respectively, by introducing cells from the Lewis lung carcinoma cell line or the PANC02 murine pancreatic adenocarcinoma cell line into an appropriate mouse strain. A mouse model of skin cancer can be
25 obtained by inducing tumors in mice by topical application of DMBA (7,12-dimethylbenz[*a*]anthracene).

Tumor Prevention

Peptides of the invention which are associated with the cancers listed above are identified and synthesized as described above.

For those mouse models in which tumors are induced by a tumorigenic
5 compound, or are produced from implanted tumor cells, the mice are initially
vaccinated with approximately 50-500 micrograms of peptide mixed with 45 ng
GM-CSF. The mice are then injected twice more with the same composition at
intervals of 2 to 4 weeks. Mice receiving GM-CSF only are used as a control.
After the third injection, the mice are inoculated with tumor cells or exposed to
10 a tumorigenic compound in a dose known to produce tumors at a high incidence
(i.e., in 75% to 90% or greater of the animals) in that particular host. The mice
are then monitored for the time to first presentation of tumor, and the sizes of
the tumors are measured every other day thereafter.

For those mouse models which spontaneously develop tumors, an initial
15 vaccination with approximately 50-500 micrograms of peptide mixed with 45 ng
GM-CSF is given at 6 weeks of age, the time at which the immune system is
developed in mice, and then at 2 to 4 week intervals thereafter. Mice receiving
GM-CSF only are used as a control. The mice are then monitored for the time
to first presentation of tumor, and the sizes of the tumors are measured every
20 other day thereafter.

Tumor Regression

Peptides of the invention which are associated with the cancers listed above are identified and synthesized as described above.

25 For those mouse models in which tumors are induced by a tumorigenic
compound, or are produced from implanted tumor cells, mice are initially
inoculated with tumor cells or exposed to the tumorigenic compound. At a time
period approximately four days prior to when tumors are expected to develop,
the mice are vaccinated with approximately 50-500 micrograms of peptide
30 mixed with 45 ng GM-CSF and injected twice more with the same composition
at intervals of 2 to 4 weeks. Mice receiving GM-CSF only are used as a control.

-61-

The mice are monitored for the time to first presentation of tumor, and the sizes of the tumors are measured every other day thereafter.

For those mouse models which spontaneously develop tumors, an initial vaccination with approximately 50-500 micrograms of peptide mixed with 45 ng GM-CSF is given at 6 weeks of age, the time at which the immune system is developed in mice, and then at 2 to 4 week intervals thereafter. Mice receiving GM-CSF only are used as a control. The mice are monitored for the time to first presentation of tumor, and the sizes of the tumors are measured every other day thereafter.

10 Tumor Prevention and Regression with Multimeric Peptides

The experiments described above for demonstrating tumor prevention and regression in mouse models of cancer are repeated using dimeric, trimeric, tetrameric, pentameric or hexameric peptides of the invention which are associated with the particular cancer.

15

Tumor Prevention and Regression with Peptide Admixtures

The experiments described above for demonstrating tumor prevention and regression in mouse models of cancer are repeated using admixtures of two, three, four, five or six peptides of the invention which are associated with the particular cancer.

20

Example 8 - Prevention or Regression of Tumors Formed by Syngeneic Mouse Tumor Cells Expressing Human Alternative Splice Forms

Peptides of the invention of human origin associated with a cancer, as listed in Table 1, are identified and synthesized as described above. Expression vectors capable of expressing some or all of the human alternative splice form sequences from which the vaccinating peptides are derived are constructed according to standard techniques.

25

Syngeneic Mouse Tumor Model

30 An NIH-3T3 cell line is transfected with the SV40 T antigen to render it tumorigenic. Alternatively, an NIH-3T3 cell line is rendered tumorigenic by

-62-

being maintained continuously in cell culture for 30-40 passages, and tumorigenic cells which spontaneously arise are isolated and propagated.

The tumorigenic NIH-3T3 cells are transfected with the plasmid expression vector described above, and cells expressing the human alternative splice form sequences are selected and propagated by standard techniques. Tumorigenic NIH-3T3 cells expressing the human alternative splice form sequences are injected into NIH Swiss mice, and tumors are allowed to form. The tumors are excised from some animals and propagated in cell culture, where the cells maintain their ability to form tumors in animals. Expression of the human sequences is confirmed. These cells are then used in subsequent tumor prevention/regression experiments.

Tumor Prevention

NIH Swiss mice are initially vaccinated with approximately 50-500 micrograms of one or more peptides of the invention of human origin described above, mixed with 45 ng GM-CSF. The mice are then injected twice more with the same composition at intervals of 2 to 4 weeks. Mice receiving GM-CSF only are used as a control. After the third injection, the mice are inoculated with syngeneic tumor cells expressing human alternative splice form sequences, as described above. The mice are then monitored for the time to first presentation of tumor, and the sizes of the tumors are measured every other day thereafter.

Tumor Regression

NIH Swiss mice are initially inoculated with syngeneic tumor cells expressing the human alternative splice form, as described above. On the fourth day post inoculation, the mice are vaccinated with approximately 50-500 micrograms of one or more peptides of the invention of human origin described above, mixed with 45 ng GM-CSF. The mice are injected twice more with the same composition at intervals of 2 to 4 weeks. Mice receiving GM-CSF only are used as a control. The mice are monitored for the time to first presentation of tumor, and the sizes of the tumors are measured every other day thereafter.

30

All documents referred to herein are incorporated by reference in their entirety. While the present invention has been described in connection with the preferred embodiments and the various figures, it is to be understood that other similar embodiments may be used or modifications and additions made to the described embodiments for performing the same function of the present invention without deviating therefrom. Therefore, the present invention should not be limited to any single embodiment, but rather should be construed in breadth and scope in accordance with the recitation of the appended claims.

5

I claim:

1. A peptide comprising an amino acid sequence unique to an alternative splice form.
2. The peptide of claim 1, wherein the peptide is from 4 to 50 amino acids in length.
3. The peptide of claim 2, wherein the peptide is from 7 to 15 amino acids in length.
4. The peptide of claim 2, wherein the peptide is 8 or 9 amino acids in length.
5. The peptide of claim 1, wherein the peptide contains at least one normal amino acid sequence of 7 contiguous amino acids or less.
6. The peptide of claim 5, wherein the peptide contains at least one normal amino acid sequence of 6, 5, 4, 3, 2 or 1 contiguous amino acids.
7. The peptide of claim 5, wherein the peptide contains at least one normal amino acid sequence of 5 or 6 contiguous amino acids.
8. The peptide of claim 5, wherein the peptide contains a splice junction, a first normal amino acid sequence of 7 contiguous amino acids or less and a second normal amino acid sequence of 7 contiguous amino acids or less, wherein the first and second normal amino acid sequences flank the alternative splice junction.
9. The peptide of claim 8, wherein the first normal amino acid sequence has 6, 5, 4, 3, 2 or 1 contiguous amino acids and the second normal amino acid sequence has 6, 5, 4, 3, 2 or 1 contiguous amino acids.

10. The peptide of claim 8, wherein the first and second normal amino acid sequences each have 3 or 4 contiguous amino acids.

11. The peptide of claim 1, wherein the peptide contains at least one modification for coupling the peptide to a polyvalent platform or to another protein.

12. The peptide of claim 11, wherein the modification comprises additional amino acids added to the C- or N-terminus of the peptide.

13. The peptide of claim 12, wherein the amino acids added to the C- or N-terminus of the peptide are selected from the group consisting of tyrosine, cysteine, lysine, glutamic acid and aspartic acid.

14. The peptide of claim 13, wherein the amino acid added to the C- or N-terminus of the peptide is cysteine.

15. The peptide of claim 11, wherein the modification is selected from the group consisting of introduction of coupling sites by terminal-NH₂ acylation; thioglycolic acid amidation; terminal-carboxy amidation; and biotinylation.

16. The peptide of claim 1, which contains at least one D-amino acid.

17. The peptide of claim 1, selected from the group consisting of SEQ ID NO: 73; SEQ ID NO: 75; SEQ ID NO: 77; and SEQ ID NO: 79.

18. A multimer comprising two or more peptides of claim 1.

19. The multimer of claim 18, wherein the multimer is a homomultimer.

20. The multimer of claim 19 which is selected from the group consisting of a dimer; trimer; tetramer; pentamer; and hexamer.

21. The multimer of claim 20, wherein the multimer is a dimer.

22. The dimer of claim 21 comprising two peptides of SEQ ID NO: 81.

23. The multimer of claim 18, wherein the multimer is a heteromultimer.

24. The multimer of claim 23 which is selected from the group consisting of a dimer; trimer; tetramer; pentamer; and hexamer.

25. A method of identifying immunogenic peptides for treating a subject who has, or is at risk for having, a disease or condition in which diseased or abnormal cells produce at least one alternative splice form, which alternative splice form is substantially absent from normal cells, comprising the steps of:

- 1) identifying at least one mRNA which encodes for the at least one alternative splice form;
- 2) determining at least a partial amino acid sequence of the at least one alternative splice form; and
- 3) generating at least one peptide comprising an amino acid sequence which is unique to the alternative splice form.

26. A method of treating a subject who has, or is at risk for having, a disease or condition in which diseased or abnormal cells produce at least one alternative splice form, which alternative splice form is substantially absent from normal cells, comprising administering an effective amount of at least one peptide of claim 1 to the subject such that an immune response is generated against the diseased or abnormal cells.

27. The method of claim 26, wherein the immune response is an MHC HLA-class I or class II restricted cytotoxic T lymphocyte response, or an antibody response.

28. The method of claim 27, wherein the cytotoxic T lymphocyte response is a CD8⁺ T lymphocyte response wherein CD8⁺, MHC class I-restricted T lymphocytes are activated.

29. The method of claim 27, wherein the cytotoxic T lymphocyte response is a CD4⁺ T lymphocyte response wherein CD4⁺, MHC class II-restricted T lymphocytes are activated.

30. The method of claim 26, wherein the subject is a human being.

31. The method of claim 26, wherein the at least one peptide is selected from the group consisting of SEQ ID NO: 3; SEQ ID NO: 5; SEQ ID NO: 8; SEQ ID NO: 10; SEQ ID NO: 12; SEQ ID NO: 26; SEQ ID NO: 29; SEQ ID NO: 35; SEQ ID NO: 36; SEQ ID NO: 38; SEQ ID NO: 39; SEQ ID NO: 73; SEQ ID NO: 75; SEQ ID NO: 77; and SEQ ID NO: 79.

32. The method of claim 26, wherein the immune response generated against the diseased or abnormal cells is enhanced by modifying the peptide to prior to administering the peptide to the subject.

33. The method of claim 32, wherein the peptide is modified by increasing the hydrophobicity of the peptide N-terminus.

34. The method of claim 32, wherein the peptide is modified with at least one amino acid insertion, deletion, or substitution which increases binding affinity of the peptide to an MHC molecule.

35. The method of claim 32, wherein the stability of the peptide is increased by modifying the peptide prior to administering the peptide to the subject.

36. The method of claim 35, wherein the peptide is modified by capping the peptide with a D-amino acid, replacing at least one L-amino acid of the peptide with a D-amino acid, or reversing the amino acid sequence of the peptide and replacing at least one L-amino acid with a D-amino acid.

37. The method of claim 26, wherein two or more peptides are administered to the subject.

38. The method of claim 37, wherein the two or more peptides comprise a multimer.

39. The method of claim 38, wherein the multimer is a homomultimer.

40. The method of claim 39, wherein the multimer is selected from the group consisting of a dimer; trimer; tetramer; pentamer; and hexamer.

41. The method of claim 40, wherein the multimer is a dimer.

42. The method of claim 41 comprising two peptides of SEQ ID NO: 81.

43. The method of claim 38, wherein the multimer is a heteromultimer.

44. The method of claim 43, wherein the multimer is selected from the group consisting of a dimer; trimer; tetramer; pentamer; and hexamer.

45. The method of claim 37, wherein the two or more peptides comprise an admixture.

46. The method of claim 37, wherein the two or more peptides comprise overlapping epitopes from one or more alternative splice forms.

47. The method of claim 26, wherein the at least one peptide is administered to the subject in combination with peptides that present T-helper cell epitopes.

48. The method of claim 26, wherein the at least one peptide is administered to the subject in combination with at least one component that primes cytotoxic T lymphocytes.

49. The method of claim 48, wherein the at least one component that primes cytotoxic T lymphocytes comprises tripalmitoyl-S-glycerylcysteinyl-serine (P₃CSS).

50. The method of claim 26, wherein the effective amount is about 1 microgram to about 2,000 mg of the at least one peptide per 70 kg of subject.

51. The method of claim 26, wherein the effective amount is about 1 microgram to about 500 mg of the at least one peptide per 70 kg of subject.

52. The method of claim 26, wherein the effective amount is about 10 micrograms to about 200 mg of the at least one peptide per 70 kg of subject.

53. The method of claim 26, wherein the effective amount is about 50 micrograms to about 100 mg of the at least one peptide per 70 kg of subject.

54. The method of claim 26, wherein the effective amount of the at least one peptide is administered in a single dose.

55. The method of claim 26, wherein the effective amount of the at least one peptide is administered in multiple doses.

56. The method of claim 26, wherein the effective amount of the at least one peptide is administered enterally.

57. The method of claim 56, wherein the enteral route of administration is selected from the group consisting of oral; rectal; and intranasal.

58. The method of claim 26, wherein the effective amount of the at least one peptide is administered parenterally.

59. The method of claim 58, wherein the parenteral route of administration is selected from the group consisting of intravenous; intramuscular; intraarterial; intraperitoneal; intravaginal; intravesical; intradermal; intrapulmonary; inhalation; topical; subcutaneous; and instillation into the body.

60. The method of claim 26, wherein the at least one peptide is administered to a subject in combination with a carrier or adjuvant.

61. The method of claim 60, wherein the carrier is selected from the group consisting of keyhole limpet hemocyanin; thyroglobulin; albumins; tetanus toxoid; and polyamino acids.

62. The method of claim 60, wherein the adjuvant is selected from the group consisting of complete Freund's adjuvant; incomplete Freund's adjuvant; aluminum phosphate; aluminum hydroxide; polylecithins; emulsified oils; and alum.

-71-

63. The method of claim 26, wherein the at least one peptide is administered to a subject in combination with an immunostimulatory compound.

64. The method of claim 63, wherein the immunostimulatory compound is selected from the group consisting of cytokines and haptens.

65. The method of claim 64, where the cytokines are selected from the group consisting of GM-CSF; IL-12; IL-2; IL-4; IL-1alpha; and IL-18.

66. The method of claim 26, wherein the effective amount of the at least one peptide is administered to a subject by expression of at least one nucleic acid sequence encoding the at least one peptide in cells of the subject

67. The method of claim 66, wherein cells of the subject are infected with an attenuated viral host comprising the at least one nucleic acid.

68. The method of claim 67, wherein the attenuated viral host is vaccinia virus.

69. The method of claim 66, wherein a bacterial host comprising the at least one nucleic acid is introduced into a subject.

70. The method of claim 69, wherein the bacterial host is selected from the group consisting of BCG; *Salmonella typhi*; and *Listeria monocytogenes*.

71. The method of claim 66, wherein a yeast host comprising the at least one nucleic acid is introduced into a subject.

72. The method of claim 71, wherein the yeast host is *Saccharomyces cerevisiae* or *Schizosaccharomyces pombe*.

73. The method of claim 26, wherein the disease or condition is selected from the group consisting of stress; cancer; diseases or conditions of the immune system; metabolic disorders; connective tissue disorders; disorders of the arteries; inherited red cell membrane disorders; thyroid hormone repression; endometrial hyperplasia; Alzheimer's disease; and alcoholism.

74. The method of claim 73, wherein the cancer is selected from the group consisting of acute promyelocytic leukemia; acute lymphoblastic leukemia; myeloblastic leukemia; uterine cancer; thyroid cancer; gastrointestinal cancer; dysplastic and neoplastic cervical epithelium; melanoma; endometrial cancer; teratocarcinoma; colon cancer; desmoplastic round cell tumors; gastric cancer; breast cancer, ovarian cancer, prostate cancer, lung cancer, skin cancer, lymphoma, bladder cancer, and pancreatic cancer.

75. The method of claim 26, wherein the disease or disorder is selected from the group consisting of allergic response; x-linked agammaglobulinemia; immunity/inflammation; systemic lupus erythematosus; Goodpasture disease; phenylketonuria; non-insulin dependent diabetes; osteogenesis imperfecta; atherosclerosis; and hereditary elliptocytosis.

76. The method of claim 26, wherein the alternative splice form is produced from the CD44 gene; the estrogen receptor gene; or the FHIT gene.

-73-

77. The method of claim 26, wherein administering the at least one peptide to a subject comprises the steps of:

- 1) removing immune system effector cells from a subject;
- 2) maintaining the immune system effector cells in culture outside the body of the subject;
- 3) optionally enriching the immune system effector cells for a particular immune system effector cell type;
- 4) treating the cultured immune system effector cells with the at least one peptide;
- 5) optionally examining a portion of the treated immune system effector cells to confirm the presence of the at least one peptide within the cells and
- 6) reintroducing the treated immune system effector cells into the subject.

78. The method of claim 77, wherein the immune system effector cells are selected from the group consisting of dendritic cells; lymphokine-activated killer cells; natural killer cells; T-cells; macrophages; and combinations thereof.

79. The method of claim 77, wherein treatment of the immune system effector cells with the at least one peptide comprises direct exposure of the cells to the at least one peptide.

80. The method of claim 77, wherein treatment of the immune system effector cells with the at least one peptide comprises introduction of at least one nucleic acid encoding the at least one peptide into the cells.

81. The method of claim 77, wherein the treated immune system effector cells are reintroduced into the subject by intravenous infusion or direct injection into the bone marrow.

82. The method of claim 77, wherein about 10^5 to about 10^8 treated immune system effector cells per kilogram of subject body weight are reintroduced into the subject.

83. A method of preventing or delaying the onset of tumor development in a subject at risk for having a tumor in which tumor cells produce at least one alternative splice form, which alternative splice form is substantially absent from non-tumor cells, comprising administering to a subject an effective amount of at least one peptide of claim 1, such that an immune response is generated against the tumor cells.

84. The method of claim 83, wherein the tumor derives from a cancer selected from the group consisting of a uterine cancer; thyroid cancer; gastrointestinal cancer; dysplastic and neoplastic cervical epithelium; melanoma; endometrial cancer; teratocarcinoma; colon cancer; desmoplastic round cell tumors; gastric cancer; breast cancer, ovarian cancer, prostate cancer, lung cancer, skin cancer, lymphoma, bladder cancer, and pancreatic cancer.

85. The method of claim 83, wherein the subject is a human being.

86. The method of claim 83, wherein two or more peptides are administered to the subject.

87. The method of claim 86, wherein the two or more peptides comprise a multimer.

88. The method of claim 87, wherein the multimer is selected from the group consisting of a dimer; trimer; tetramer; pentamer; and hexamer.

89. The method of claim 87, wherein the multimer is a dimer and comprises two peptides of SEQ ID NO: 81.

90. The method of claim 86, wherein the two or more peptides comprise an admixture.

91. The method of claim 83, wherein the at least one peptide is selected from the group consisting of SEQ ID NO: 8; SEQ ID NO: 10; SEQ ID NO: 26; SEQ ID NO: 29; SEQ ID NO: 73; SEQ ID NO: 75; SEQ ID NO: 77; and SEQ ID NO: 79.

92. A method of regressing a tumor in a subject having a tumor in which tumor cells produce at least one alternative splice form, which alternative splice form is substantially absent from non-tumor cells, comprising administering to a subject an effective amount of at least one peptide of claim 1, such that an immune response is generated against the tumor cells.

93. The method of claim 92, wherein the tumor derives from a cancer selected from the group consisting of a uterine cancer; thyroid cancer; gastrointestinal cancer; dysplastic and neoplastic cervical epithelium; melanoma; endometrial cancer; teratocarcinoma; colon cancer; desmoplastic round cell tumors; gastric cancer; breast cancer, ovarian cancer, prostate cancer, lung cancer, skin cancer, lymphoma, bladder cancer, and pancreatic cancer.

94. The method of claim 92, wherein the subject is a human being.

95. The method of claim 88, wherein two or more peptides are administered to the subject.

96. The method of claim 95, wherein the two or more peptides comprise a multimer.

97. The method of claim 96, wherein the multimer is selected from the group consisting of a dimer; trimer; tetramer; pentamer; and hexamer.

98. The method of claim 97, wherein the multimer is a dimer and comprises two peptides of SEQ ID NO: 81.

99. The method of claim 95, wherein the two or more peptides comprise an admixture.

100. The method of claim 92, wherein the at least one peptide is selected from the group consisting of SEQ ID NO: 8; SEQ ID NO: 10; SEQ ID NO: 26; SEQ ID NO: 29; SEQ ID NO: 73; SEQ ID NO: 75; SEQ ID NO: 77; and SEQ ID NO: 79.

101. A method for identifying peptides which induce MHC-restricted cytotoxic T lymphocyte responses in a subject, comprising the steps of:

- 1) obtaining peripheral blood lymphocytes;
- 2) exposing the peripheral blood lymphocytes to at least one peptide of claim 1 such that the peripheral blood lymphocytes are stimulated;
- 3) incubating the stimulated peripheral blood lymphocytes with target cells that either endogenously synthesize the alternative splice form from which the peptide is derived or are pulsed with the peptide; and
- 4) detecting lysis of the target cells.

102. The method of claim 101, wherein the target cells are autologously labeled and the detection of target cell lysis comprises measuring release of the autologous label from the lysed target cells.

103. The method of claim 101, wherein the detection of target cell lysis comprises detection of at least one cytokine released from activated peripheral blood lymphocytes upon lysis of the target cells.

104. The method of claim 102, wherein the at least one cytokine is interferon-gamma.

-77-

105. An antibody or antibody fragment that binds to specific epitopes on a peptide selected from the group consisting of SEQ ID NO: 73; SEQ ID NO: 75; SEQ ID NO: 77; and SEQ ID NO: 79.

106. The antibody of claim 105 which is humanized.

107. The antibody of claim 105 which is a polyclonal antibody.

108. The antibody of claim 105 which is a monoclonal antibody.

109. A hybridoma producing the monoclonal antibody of claim 108.

110. A method of treating a subject having, or at risk for having, a disease or condition in which diseased or abnormal cells produce at least one alternative splice form, which alternative splice form is substantially absent from normal cells, comprising administering to a subject an effective amount of at least one antibody specific to an amino acid sequence unique to the alternative splice form, such that one or more clinical symptoms in the subject are ameliorated or the number of diseased or abnormal cells in the subject is reduced.

111. The method of claim 110, wherein the at least one antibody is a monoclonal antibody.

112. The method of claim 110, wherein the at least one antibody is humanized.

113. The method of claim 111, wherein the monoclonal antibody binds to specific epitopes on a peptide selected from the group consisting of SEQ ID NO: 3; SEQ ID NO: 5; SEQ ID NO: 8; SEQ ID NO: 10; SEQ ID NO: 12; SEQ ID NO: 26; SEQ ID NO: 29; SEQ ID NO: 35; SEQ ID NO: 36; SEQ

-78-

ID NO: 38; SEQ ID NO: 39; SEQ ID NO: 73; SEQ ID NO: 75; SEQ ID NO: 77;
and SEQ ID NO: 79.

114. The method of claim 110, wherein the effective amount is about 0.1 mg/kg to about 100 mg/kg of body weight.

115. The method of claim 110, wherein the effective amount is about 50 mg/kg to about 100 mg/kg of body weight.

116. The method of claim 110, wherein the effective amount is about 10 mg/kg to about 20 mg/kg of body weight.

117. A pharmaceutical composition comprising at least one peptide of claim 1 and a pharmaceutically acceptable carrier.

118. The pharmaceutical composition of claim 117 which is encapsulated in a liposome.

1/5




FIG. 1B

FIG. 1A

2/5






FIG. 2



4/5




FIG. 4A




FIG. 4B




FIG. 4C

5/5




FIG. 5

<110> Thomas Jefferson University
Albert J. Wong

<120> Alternative Splice Forms of Proteins as
Basis for Multiple Therapeutic Modalities

<130> 8321-81 PC

<150> US 60/293,791

<151> 2001-05-25

<160> 82

<170> FastSEQ for Windows Version 4.0

<210> 1

<211> 271

<212> PRT

<213> Homo Sapiens

<400> 1

```

Met Phe Gly Leu Lys Arg Asn Ala Val Ile Gly Leu Asn Leu Tyr Cys
 1      5      10
Gly Gly Ala Gly Leu Gly Ala Gly Ser Gly Gly Ala Thr Arg Pro Gly
 20      25
Gly Arg Leu Leu Ala Thr Glu Lys Glu Ala Ser Ala Arg Arg Glu Ile
 35      40      45
Gly Gly Gly Glu Ala Gly Ala Val Ile Gly Gly Ser Ala Gly Ala Ser
 50      55      60
Pro Pro Ser Thr Leu Thr Pro Asp Ser Arg Arg Val Ala Arg Pro Pro
 65      70      75
Pro Ile Gly Ala Glu Val Pro Asp Val Thr Ala Thr Pro Ala Arg Leu
 85      90      95
Leu Phe Phe Ala Pro Thr Arg Arg Ala Ala Pro Leu Glu Glu Met Glu
100      105      110
Ala Pro Ala Ala Asp Ala Ile Met Ser Pro Glu Glu Glu Leu Asp Gly
115      120      125
Tyr Glu Pro Glu Pro Leu Gly Lys Arg Pro Ala Val Leu Pro Leu Leu
130      135      140
Glu Leu Val Gly Glu Ser Gly Asn Asn Thr Ser Thr Asp Gly Ser Leu
145      150      155
Pro Ser Thr Pro Pro Ala Glu Glu Glu Glu Asp Glu Leu Tyr Arg
165      170      175
Gln Ser Leu Glu Ile Ile Ser Arg Tyr Leu Arg Glu Gln Ala Thr Gly
180      185      190
Ala Lys Asp Thr Lys Pro Met Gly Arg Ser Gly Ala Thr Ser Arg Lys
195      200      205
Ala Leu Glu Thr Leu Arg Arg Val Gly Asp Gly Val Gln Arg Asn His
210      215      220
Glu Thr Ala Phe Gln Gly Trp Val Cys Gly Val Leu Pro Cys Arg Gly
225      230      235
Pro Arg Arg Trp His Gln Glu Cys Ala Ala Gly Phe Cys Arg Cys Cys
245      250      255
Trp Ser Arg Ser Trp Phe Gly Ile Ser Asn Lys Ile Ala Leu Leu
260      265      270

```

<210> 2

<211> 18

<212> PRT

<213> Homo sapiens

<400> 2

Arg Asn His Glu Thr Ala Phe Gln Gly Trp Val Cys Gly Val Leu Pro
 1 5 10 15
 Cys Arg

<210> 3

<211> 61

<212> PRT

<213> Homo Sapiens

<400> 3

Asn Ser Asn His Val Ala Ser Gly Ala Pro Val Cys His Asn Pro Asn
 1 5 10 15
 Leu Pro Ser Trp Gln Gly Ala Leu Gly Pro Tyr Gly Val Val Val Leu
 20 25 30
 Ala Pro Asp Thr Trp Leu Ser Ser Leu Arg Leu Ser Ser Pro Gly
 35 40 45
 Val Glu Gly Arg Ser Cys Ser Ala Arg Glu Thr Gln Ala
 50 55 60

<210> 4

<211> 682

<212> PRT

<213> Homo Sapiens

<400> 4

Met Ala Gly Gly Pro Gly Pro Gly Glu Pro Ala Ala Pro Gly Ala Gln
 1 5 10 15
 His Phe Leu Tyr Glu Val Pro Pro Trp Val Met Cys Arg Phe Tyr Lys
 20 25 30
 Val Met Asp Ala Leu Glu Pro Ala Asp Trp Cys Gln Phe Ala Ala Leu
 35 40 45
 Ile Val Arg Asp Gln Thr Glu Leu Arg Leu Cys Glu Arg Ser Gly Gln
 50 55 60
 Arg Thr Ala Ser Val Leu Trp Pro Trp Ile Asn Arg Asn Ala Arg Val
 65 70 75 80
 Ala Asp Leu Val His Ile Leu Thr His Leu Gln Leu Leu Arg Ala Arg
 85 90 95
 Asp Ile Ile Thr Ala Trp His Pro Pro Ala Pro Leu Pro Ser Pro Gly
 100 105 110
 Thr Thr Ala Pro Arg Pro Ser Ser Ile Pro Ala Pro Ala Glu Ala Glu
 115 120 125
 Ala Trp Ser Pro Arg Lys Leu Pro Ser Ser Ala Ser Thr Phe Leu Ser
 130 135 140
 Pro Ala Phe Pro Gly Ser Gln Thr His Ser Gly Pro Glu Leu Gly Leu
 145 150 155 160
 Val Pro Ser Pro Ala Ser Leu Trp Pro Pro Pro Ser Pro Ala Pro
 165 170 175
 Ser Ser Thr Lys Pro Gly Pro Glu Ser Ser Val Ser Leu Leu Gln Gly
 180 185 190
 Ala Arg Pro Phe Pro Phe Cys Trp Pro Leu Cys Glu Ile Ser Arg Gly
 195 200 205
 Thr His Asn Phe Ser Glu Glu Leu Lys Ile Gly Glu Gly Gly Phe Gly
 210 215 220
 Cys Val Tyr Arg Ala Val Met Arg Asn Thr Val Tyr Ala Val Lys Arg
 225 230 235 240
 Leu Lys Glu Asn Ala Asp Leu Glu Trp Thr Ala Val Lys Gln Ser Phe
 245 250 255
 Leu Thr Glu Val Glu Gln Leu Ser Arg Phe Arg His Pro Asn Ile Val
 260 265 270
 Asp Phe Ala Gly Tyr Cys Ala Gln Asn Gly Phe Tyr Cys Leu Val Tyr
 275 280 285
 Gly Phe Leu Pro Asn Gly Ser Leu Glu Asp Arg Leu His Cys Gln Thr
 290 295 300

Gln Ala Cys Pro Pro Leu Ser Trp Pro Gln Arg Leu Asp Ile Leu Leu
 305 310 315 320
 Gly Thr Ala Arg Ala Ile Gln Phe Leu His Gln Asp Ser Pro Ser Leu
 325 330 335
 Ile His Gly Asp Ile Lys Ser Ser Asn Val Leu Leu Asp Glu Arg Leu
 340 345 350
 Thr Pro Lys Leu Gly Asp Phe Gly Leu Ala Arg Phe Ser Arg Phe Ala
 355 360 365
 Gly Ser Ser Pro Ser Gln Ser Met Val Ala Arg Thr Gln Thr Val
 370 375 380
 Arg Gly Thr Leu Ala Tyr Leu Pro Glu Glu Tyr Ile Lys Thr Gly Arg
 385 390 395 400
 Leu Ala Val Asp Thr Asp Thr Phe Ser Phe Gly Val Val Val Leu Glu
 405 410 415
 Thr Leu Ala Gly Gln Arg Ala Val Lys Thr His Gly Ala Arg Thr Lys
 420 425 430
 Tyr Leu Lys Asp Leu Val Glu Glu Glu Ala Glu Glu Ala Gly Val Ala
 435 440 445
 Leu Arg Ser Thr Gln Ser Thr Leu Gln Ala Gly Leu Ala Ala Asp Ala
 450 455 460
 Trp Ala Ala Pro Ile Ala Met Gln Ile Tyr Lys Lys His Leu Asp Pro
 465 470 475 480
 Arg Pro Gly Pro Cys Pro Pro Glu Leu Gly Leu Gly Leu Gly Gln Leu
 485 490 495
 Ala Cys Cys Cys Leu His Arg Arg Ala Lys Arg Arg Pro Pro Met Thr
 500 505 510
 Gln Glu Asn Ser Tyr Val Ser Ser Thr Gly Arg Ala His Ser Gly Ala
 515 520 525
 Ala Pro Trp Gln Pro Leu Ala Ala Pro Ser Gly Ala Ser Ala Gln Ala
 530 535 540
 Ala Glu Gln Leu Gln Arg Gly Pro Asn Gln Pro Val Glu Ser Asp Glu
 545 550 555 560
 Ser Leu Gly Gly Leu Ser Ala Ala Leu Arg Ser Trp His Leu Thr Pro
 565 570 575
 Ser Cys Pro Leu Asp Pro Ala Pro Leu Arg Glu Ala Gly Cys Pro Gln
 580 585 590
 Gly Asp Thr Ala Gly Glu Ser Ser Trp Gly Ser Gly Pro Gly Ser Arg
 595 600 605
 Pro Thr Ala Val Glu Gly Leu Ala Leu Gly Ser Ser Ala Ser Ser Ser
 610 615 620
 Ser Glu Pro Pro Gln Ile Ile Ile Asn Pro Ala Arg Gln Lys Met Val
 625 630 635 640
 Gln Lys Leu Ala Leu Tyr Glu Asp Gly Ala Leu Asp Ser Leu Gln Leu
 645 650 655
 Leu Ser Ser Ser Ser Leu Pro Gly Leu Gly Leu Glu Gln Asp Arg Gln
 660 665 670
 Gly Pro Glu Glu Ser Asp Glu Phe Gln Ser
 675 680

<210> 5
 <211> 30
 <212> PRT
 <213> Homo sapiens

<400> 5
 Val Tyr Glu Arg Leu Glu Lys Leu Gln Ala Val Val Ala Gly Val Pro
 1 5 10 15
 Gly His Ser Glu Ala Ala Ser Cys Ile Pro Pro Ser Pro Gln
 20 25 30

<210> 6
 <211> 380
 <212> PRT
 <213> Homo Sapiens

<400> 6
 Met Ala Ala Ala Leu Gly Gln Ile Trp Ala Arg Lys Leu Leu Ser
 1 5 10 15
 Val Pro Trp Leu Leu Cys Gly Pro Arg Arg Tyr Ala Ser Ser Ser Phe
 20 25 30
 Lys Ala Ala Asp Leu Gln Leu Glu Met Thr Gln Lys Pro His Lys Lys
 35 40 45
 Pro Gly Pro Gly Glu Pro Leu Val Phe Gly Lys Thr Phe Thr Asp His
 50 55 60
 Met Leu Met Val Glu Trp Asn Asp Lys Gly Trp Gly Gln Pro Arg Ile
 65 70 75 80
 Gln Pro Phe Gln Asn Leu Thr Leu His Pro Ala Ser Ser Ser Leu His
 85 90 95
 Tyr Ser Leu Gln Leu Phe Glu Gly Met Lys Ala Phe Lys Gly Lys Asp
 100 105 110
 Gln Gln Val Arg Leu Phe Arg Pro Trp Leu Asn Met Asp Arg Met Leu
 115 120 125
 Arg Ser Ala Met Arg Leu Cys Leu Pro Ser Phe Asp Lys Leu Glu Leu
 130 135 140
 Leu Glu Cys Ile Arg Arg Leu Ile Glu Val Asp Lys Asp Trp Val Pro
 145 150 155 160
 Asp Ala Ala Gly Thr Ser Leu Tyr Val Arg Pro Val Leu Ile Gly Asn
 165 170 175
 Glu Pro Ser Leu Gly Val Ser Gln Pro Arg Arg Ala Leu Leu Phe Val
 180 185 190
 Ile Leu Cys Pro Val Gly Ala Tyr Phe Pro Gly Gly Ser Val Thr Pro
 195 200 205
 Val Ser Leu Leu Ala Asp Pro Ala Phe Ile Arg Ala Trp Val Gly Gly
 210 215 220
 Val Gly Asn Tyr Lys Leu Gly Gly Asn Tyr Gly Pro Thr Val Leu Val
 225 230 235 240
 Gln Gln Glu Ala Leu Lys Arg Gly Cys Glu Gln Val Leu Trp Leu Tyr
 245 250 255
 Gly Pro Asp His Gln Leu Thr Glu Val Gly Thr Met Asn Ile Phe Val
 260 265 270
 Tyr Trp Thr His Glu Asp Gly Val Leu Glu Leu Val Thr Pro Pro Leu
 275 280 285
 Asn Gly Val Ile Leu Pro Gly Val Val Arg Gln Ser Leu Leu Asp Met
 290 295 300
 Ala Gln Thr Trp Gly Glu Phe Arg Val Val Glu Arg Thr Ile Thr Met
 305 310 315 320
 Lys Gln Leu Leu Arg Ala Leu Glu Glu Gly Arg Val Arg Glu Val Phe
 325 330 335
 Gly Ser Gly Thr Ala Cys Gln Asn Leu His Ile Pro Thr Met Glu Asn
 340 345 350
 Gly Pro Glu Leu Ile Leu Arg Phe Gln Lys Glu Leu Lys Glu Ile Gln
 355 360 365
 Tyr Gly Ile Arg Ala His Glu Trp Met Phe Pro Val
 370 375 380

<210> 7
 <211> 612
 <212> PRT
 <213> Homo Sapiens

<400> 7
 Ala Lys Leu Gly Ala Val Tyr Thr Glu Gly Gly Phe Val Glu Gly Val
 1 5 10 15
 Asn Lys Lys Leu Gly Leu Leu Gly Asp Ser Val Asp Ile Phe Lys Gly
 20 25 30
 Ile Pro Phe Ala Ala Pro Thr Lys Ala Leu Glu Asn Pro Gln Pro His
 35 40 45
 Pro Gly Trp Gln Gly Thr Leu Lys Ala Lys Asn Phe Lys Lys Arg Cys
 50 55 60
 Leu Gln Ala Thr Ile Thr Gln Asp Ser Thr Tyr Gly Asp Glu Asp Cys
 65 70 75 80

Leu Tyr Leu Asn Ile Trp Val Pro Gln Gly Arg Lys Gln Val Ser Arg
 85 90 95
 Asp Leu Pro Val Met Ile Trp Ile Tyr Gly Gly Ala Phe Leu Met Gly
 100 105 110
 Ser Gly His Gly Ala Asn Phe Leu Asn Asn Tyr Leu Tyr Asp Gly Glu
 115 120 125
 Glu Ile Ala Thr Arg Gly Asn Val Ile Val Val Thr Phe Asn Tyr Arg
 130 135 140
 Val Gly Pro Leu Gly Phe Leu Ser Thr Gly Asp Ala Asn Leu Pro Gly
 145 150 155 160
 Asn Tyr Gly Leu Arg Asp Gln His Met Ala Ile Ala Trp Val Lys Arg
 165 170 175
 Asn Ile Ala Ala Phe Gly Gly Asp Pro Asn Asn Ile Thr Leu Phe Gly
 180 185 190
 Glu Ser Ala Gly Gly Ala Ser Val Ser Leu Gln Thr Leu Ser Pro Tyr
 195 200 205
 Asn Lys Gly Leu Ile Arg Arg Ala Ile Ser Gln Ser Gly Val Ala Leu
 210 215 220
 Ser Pro Trp Val Ile Gln Lys Asn Pro Leu Phe Trp Ala Lys Lys Val
 225 230 235 240
 Ala Glu Lys Val Gly Cys Pro Val Gly Asp Ala Ala Arg Met Ala Gln
 245 250 255
 Cys Leu Lys Val Thr Asp Pro Arg Ala Leu Thr Leu Ala Tyr Lys Val
 260 265 270
 Pro Leu Ala Gly Leu Glu Tyr Pro Met Leu His Tyr Val Gly Phe Val
 275 280 285
 Pro Val Ile Asp Gly Asp Phe Ile Pro Ala Asp Pro Ile Asn Leu Tyr
 290 295 300
 Ala Asn Ala Ala Asp Ile Asp Tyr Ile Ala Gly Thr Asn Asn Met Asp
 305 310 315 320
 Gly His Ile Phe Ala Ser Ile Asp Met Pro Ala Ile Asn Lys Gly Asn
 325 330 335
 Lys Lys Val Thr Glu Glu Asp Phe Tyr Lys Leu Val Ser Glu Phe Thr
 340 345 350
 Ile Thr Lys Gly Leu Arg Gly Ala Lys Thr Thr Phe Asp Val Tyr Thr
 355 360 365
 Glu Ser Trp Ala Gln Asp Pro Ser Gln Glu Asn Lys Lys Lys Thr Val
 370 375 380
 Val Asp Phe Glu Thr Asp Val Leu Phe Leu Val Pro Thr Glu Ile Ala
 385 390 395 400
 Leu Ala Gln His Arg Ala Asn Ala Lys Ser Ala Lys Thr Tyr Ala Tyr
 405 410 415
 Leu Phe Ser His Pro Ser Arg Met Pro Val Tyr Pro Lys Trp Val Gly
 420 425 430
 Ala Asp His Ala Asp Asp Ile Gln Tyr Val Phe Gly Lys Pro Phe Ala
 435 440 445
 Thr Pro Thr Gly Tyr Arg Pro Gln Asp Arg Thr Val Ser Lys Ala Met
 450 455 460
 Ile Ala Tyr Trp Thr Asn Phe Ala Lys Thr Gly Asp Pro Asn Met Gly
 465 470 475 480
 Asp Ser Ala Val Pro Thr His Trp Glu Pro Tyr Thr Thr Glu Asn Ser
 485 490 495
 Gly Tyr Leu Glu Ile Thr Lys Lys Met Gly Ser Ser Ser Met Lys Arg
 500 505 510
 Ser Leu Arg Thr Asn Phe Leu Arg Tyr Trp Thr Leu Thr Tyr Leu Ala
 515 520 525
 Leu Pro Thr Val Thr Asp Gln Glu Ala Thr Pro Val Pro Pro Thr Gly
 530 535 540
 Asp Ser Glu Ala Thr Pro Val Pro Pro Thr Gly Asp Ser Glu Thr Ala
 545 550 555 560
 Pro Val Pro Pro Thr Gly Asp Ser Gly Ala Pro Pro Val Pro Pro Thr
 565 570 575
 Gly Asp Ser Gly Ala Pro Pro Val Pro Thr Gly Asp Ser Glu Ala
 580 585 590
 Ala Pro Val Pro Pro Thr Gly Asp Ser Lys Glu Ala Gln Met Pro Ala
 595 600 605
 Val Ile Arg Phe

610

<210> 8
 <211> 16
 <212> PRT
 <213> Homo sapiens

<400> 8
 Ala Pro Val Pro Pro Thr Gly Asp Ser Gly Ala Pro Pro Val Pro Pro
 1 5 10 15

<210> 9
 <211> 271
 <212> PRT
 <213> Homo Sapiens

<400> 9
 Met Ala Thr Thr Gly Ala Leu Gly Asn Tyr Tyr Val Asp Ser Phe Leu
 1 5 10 15
 Leu Gly Ala Asp Ala Ala Asp Glu Leu Ser Val Gly Arg Tyr Ala Pro
 20 25 30
 Gly Thr Leu Gly Gln Pro Pro Arg Gln Ala Ala Thr Leu Ala Glu His
 35 40 45
 Pro Asp Phe Ser Pro Cys Ser Phe Gln Ser Lys Ala Thr Val Phe Val
 50 55 60
 Ala Ser Trp Asn Pro Val His Ala Ala Gly Ala Asn Ala Val Pro Ala
 65 70 75 80
 Val Tyr His His His His His His Pro Tyr Val His Pro Gln Ala Pro
 85 90 95
 Val Ala Ala Ala Ala Pro Asp Gly Arg Tyr Met Arg Ser Trp Leu Glu
 100 105 110
 Pro Thr Pro Gly Ala Leu Ser Phe Ala Gly Leu Pro Ser Arg Pro
 115 120 125
 Tyr Gly Ile Lys Pro Glu Pro Leu Ser Ala Arg Arg Gly Asp Cys Pro
 130 135 140
 Thr Leu Asp Thr His Thr Leu Ser Leu Thr Asp Tyr Ala Cys Gly Ser
 145 150 155 160
 Pro Pro Val Asp Arg Glu Lys Gln Pro Ser Glu Gly Ala Phe Ser Glu
 165 170 175
 Asn Asn Ala Glu Asn Glu Ser Gly Gly Asp Lys Pro Pro Ile Asp Pro
 180 185 190
 Asn Asn Pro Ala Ala Asn Trp Leu His Ala Arg Ser Thr Arg Lys Lys
 195 200 205
 Arg Cys Pro Tyr Thr Lys His Gln Thr Leu Glu Leu Glu Lys Glu Phe
 210 215 220
 Leu Phe Asn Met Tyr Leu Thr Arg Asp Arg Arg Tyr Glu Val Ala Arg
 225 230 235 240
 Leu Phe Asn Leu Thr Glu Arg Gln Val Lys Ile Trp Phe Gln Asn Arg
 245 250 255
 Arg Met Lys Met Lys Lys Ile Asn Lys Asp Arg Ala Lys Asp Glu
 260 265 270

<210> 10
 <211> 14
 <212> PRT
 <213> Homo sapiens

<400> 10
 Asp Lys Pro Pro Ile Asp Pro Asn Asn Pro Ala Ala Asn Trp
 1 5 10

<210> 11
 <211> 587

<212> PRT

<213> Mus musculus

<400> 11

Met Ser Trp Ala Arg Ser Arg Leu Cys Ser Thr Leu Ser Leu Ala Ala
 1 5 10 15
 Val Ser Ala Arg Gly Ala Thr Thr Glu Gly Pro Ala Arg Arg Gly Met
 20 25 30
 Ser Ala Gly Pro Ala Pro Gln Glu Pro Gly Met Glu Tyr Gln Asp Ala
 35 40 45
 Val Arg Thr Leu Asn Thr Leu Gln Thr Asn Ala Ser Tyr Leu Glu Gln
 50 55 60
 Val Lys Arg Gln Arg Ser Asp Pro Gln Ala Gln Leu Glu Ala Met Glu
 65 70 75 80
 Met Tyr Leu Ala Arg Ser Gly Leu Gln Val Glu Asp Leu Asn Arg Leu
 85 90 95
 Asn Ile Ile His Val Thr Gly Thr Lys Gly Lys Gly Ser Thr Cys Ala
 100 105 110
 Phe Thr Glu Arg Ile Leu Arg Asn Tyr Gly Leu Lys Thr Gly Phe Phe
 115 120 125
 Arg Ser Pro His Met Val Gln Val Arg Asp Arg Ile Arg Ile Asn Gly
 130 135 140
 Lys Pro Ile Ser Pro Glu Leu Phe Thr Lys His Phe Trp Cys Leu Tyr
 145 150 155 160
 Asn Gln Leu Glu Glu Phe Lys Asp Asp Ser His Val Ser Met Pro Ser
 165 170 175
 Tyr Phe Arg Phe Leu Thr Leu Met Ala Phe His Val Phe Leu Gln Glu
 180 185 190
 Lys Val Asp Leu Ala Val Val Glu Val Gly Ile Gly Gly Ala Phe Asp
 195 200 205
 Cys Thr Asn Ile Ile Arg Lys Pro Val Val Cys Gly Val Ser Ser Leu
 210 215 220
 Gly Ile Asp His Thr Ser Leu Leu Gly Asp Thr Val Glu Lys Ile Ala
 225 230 235 240
 Trp Gln Lys Gly Gly Ile Phe Lys Pro Gly Val Pro Ala Phe Thr Val
 245 250 255
 Val Gln Pro Glu Gly Pro Leu Ala Val Leu Arg Asp Arg Ala Gln Gln
 260 265 270
 Ile Gly Cys Pro Leu Tyr Leu Cys Pro Pro Leu Glu Ala Leu Glu Glu
 275 280 285
 Val Gly Leu Pro Leu Ser Leu Gly Leu Glu Gly Ala His Gln Arg Ser
 290 295 300
 Asn Ala Ala Leu Ala Leu Gln Leu Ala His Cys Trp Leu Glu Arg Gln
 305 310 315 320
 Asp His Gln Asp Ile Gln Glu Leu Lys Val Ser Arg Pro Ser Ile Arg
 325 330 335
 Trp Gln Leu Pro Leu Ala Pro Val Phe Arg Pro Thr Pro His Met Arg
 340 345 350
 Arg Gly Leu Arg Asp Thr Val Trp Pro Gly Arg Thr Gln Ile Leu Gln
 355 360 365
 Arg Gly Pro Leu Thr Trp Tyr Leu Asp Gly Ala His Thr Thr Ser Ser
 370 375 380
 Val Gln Ala Cys Val His Trp Tyr Arg Gln Ser Leu Glu Arg Ser Lys
 385 390 395 400
 Arg Thr Asp Gly Gly Ser Glu Val His Ile Leu Leu Phe Asn Ser Thr
 405 410 415
 Gly Asp Arg Asp Ser Ala Ala Leu Leu Lys Leu Leu Gln Pro Cys Gln
 420 425 430
 Phe Asp Tyr Ala Val Phe Cys Pro Asn Val Thr Glu Val Ser Ser Ile
 435 440 445
 Gly Asn Ala Asp Gln Gln Asn Phe Thr Val Thr Leu Asp Gln Val Leu
 450 455 460
 Leu Arg Cys Leu Gln His Gln Gln His Trp Asn Gly Leu Ala Glu Lys
 465 470 475 480
 Gln Ala Ser Ser Asn Leu Trp Ser Ser Cys Gly Pro Asp Pro Ala Gly
 485 490 495
 Pro Gly Ser Leu Leu Leu Ala Pro His Pro Pro Gln Pro Thr Arg Thr

```

          500          505          510
Ser Ser Leu Val Phe Ser Cys Ile Ser His Ala Leu Leu Trp Ile Ser
      515          520          525
Gln Gly Arg Asp Pro Ile Phe Gln Pro Gln Ser Leu Pro Arg Asn Leu
      530          535          540
Leu Asn His Pro Thr Ala Asn Ser Gly Ala Ser Ile Leu Arg Glu Ala
545          550          555          560
Ala Ala Ile His Val Leu Val Thr Gly Ser Leu His Leu Val Gly Gly
      565          570          575
Val Leu Lys Leu Leu Asp Pro Ser Met Ser Gln
      580          585

```

<210> 12
 <211> 18
 <212> PRT
 <213> Mus musculus

```

<400> 12
Ala Val Ser Ala Arg Gly Ala Thr Thr Glu Gly Pro Ala Arg Arg Gly
  1          5          10          15
Met Ser

```

<210> 13
 <211> 237
 <212> DNA
 <213> Homo Sapiens

<220>
 <221> CDS
 <222> (1)...(237)

```

<400> 13
gac aga atc cct gct acc aat agg aat gat gtc aca ggt gga aga aga      48
Asp Arg Ile Pro Ala Thr Asn Arg Asn Asp Val Thr Gly Gly Arg Arg
  1          5          10          15

gac cca aat cat tct gaa ggc tca act act tta ctg gaa ggt tat acc      96
Asp Pro Asn His Ser Glu Gly Ser Thr Thr Leu Leu Glu Gly Tyr Thr
          20          25          30

tct cat tac cca cac acg aag gaa agc agg acc ttc atc cca gtg acc      144
Ser His Tyr Pro His Thr Lys Glu Ser Arg Thr Phe Ile Pro Val Thr
          35          40          45

tca gct aag act ggg tcc ttt gga gtt act gca gtt act gtt gga gat      192
Ser Ala Lys Thr Gly Ser Phe Gly Val Thr Ala Val Thr Val Gly Asp
      50          55          60

tcc aac tct aat gtc aat cgt tcc tta tca gga gac caa gac aca      237
Ser Asn Ser Asn Val Asn Arg Ser Leu Ser Gly Asp Gln Asp Thr
      65          70          75

```

<210> 14
 <211> 79
 <212> PRT
 <213> Homo Sapiens

```

<400> 14
Asp Arg Ile Pro Ala Thr Asn Arg Asn Asp Val Thr Gly Gly Arg Arg
  1          5          10          15
Asp Pro Asn His Ser Glu Gly Ser Thr Thr Leu Leu Glu Gly Tyr Thr
      20          25          30

```

Ser His Tyr Pro His Thr Lys Glu Ser Arg Thr Phe Ile Pro Val Thr
 35 40 45
 Ser Ala Lys Thr Gly Ser Phe Gly Val Thr Ala Val Thr Val Gly Asp
 50 55 60
 Ser Asn Ser Asn Val Asn Arg Ser Leu Ser Gly Asp Gln Asp Thr
 65 70 75

<210> 15
 <211> 327
 <212> DNA
 <213> Homo Sapiens

<220>
 <221> CDS
 <222> (1)...(327)

<400> 15
 gac aga atc cct gct acc aag cag agt aat tct cag agc ttc tct aca 48
 Asp Arg Ile Pro Ala Thr Lys Gln Ser Asn Ser Gln Ser Phe Ser Thr
 1 5 10 15
 tca cat gaa ggc ttg gaa gaa gat aaa gac cat cca aca act tct act 96
 Ser His Glu Gly Leu Glu Glu Asp Lys Asp His Pro Thr Thr Ser Thr
 20 25 30
 ctg aca tca agc aat agg aat gat gtc aca ggt gga aga aga gac cca 144
 Leu Thr Ser Ser Asn Arg Asn Asp Val Thr Gly Gly Arg Arg Asp Pro
 35 40 45
 aat cat tct gaa ggc tca act act tta ctg gaa ggt tat acc tct cat 192
 Asn His Ser Glu Gly Ser Thr Thr Leu Leu Glu Gly Tyr Thr Ser His
 50 55 60
 tac cca cac acg aag gaa agc agg acc ttc atc cca gtg acc tca gct 240
 Tyr Pro His Thr Lys Glu Ser Arg Thr Phe Ile Pro Val Thr Ser Ala
 65 70 75 80
 aag act ggg tcc ttt gga gtt act gca gtt act gtt gga gat tcc aac 288
 Lys Thr Gly Ser Phe Gly Val Thr Ala Val Thr Val Gly Asp Ser Asn
 85 90 95
 tct aat gtc aat cgt tcc tta tca gga gac caa gac aca 327
 Ser Asn Val Asn Arg Ser Leu Ser Gly Asp Gln Asp Thr
 100 105

<210> 16
 <211> 109
 <212> PRT
 <213> Homo Sapiens

<400> 16
 Asp Arg Ile Pro Ala Thr Lys Gln Ser Asn Ser Gln Ser Phe Ser Thr
 1 5 10 15
 Ser His Glu Gly Leu Glu Glu Asp Lys Asp His Pro Thr Thr Ser Thr
 20 25 30
 Leu Thr Ser Ser Asn Arg Asn Asp Val Thr Gly Gly Arg Arg Asp Pro
 35 40 45
 Asn His Ser Glu Gly Ser Thr Thr Leu Leu Glu Gly Tyr Thr Ser His
 50 55 60
 Tyr Pro His Thr Lys Glu Ser Arg Thr Phe Ile Pro Val Thr Ser Ala
 65 70 75 80
 Lys Thr Gly Ser Phe Gly Val Thr Ala Val Thr Val Gly Asp Ser Asn
 85 90 95
 Ser Asn Val Asn Arg Ser Leu Ser Gly Asp Gln Asp Thr
 100 105

<210> 17
 <211> 429
 <212> DNA
 <213> Homo Sapiens

<220>
 <221> CDS
 <222> (1)...(429)

<400> 17
 gac aga atc cct gct acc aat atg gac tcc agt cat agt aca acg ctt 48
 Asp Arg Ile Pro Ala Thr Asn Met Asp Ser Ser His Ser Thr Thr Leu
 1 5 10 15
 cag cct act gca aat cca aac aca ggt ttg gtg gaa gat ttg gac agg 96
 Gln Pro Thr Ala Asn Pro Asn Thr Gly Leu Val Glu Asp Leu Asp Arg
 20 25 30
 aca gga cct ctt tca atg aca acg cag cag agt aat tct cag agc ttc 144
 Thr Gly Pro Leu Ser Met Thr Thr Gln Gln Ser Asn Ser Gln Ser Phe
 35 40 45
 tct aca tca cat gaa ggc ttg gaa gaa gat aaa gac cat cca aca act 192
 Ser Thr Ser His Glu Gly Leu Glu Glu Asp Lys Asp His Pro Thr Thr
 50 55 60
 tct act ctg aca tca agc aat agg aat gat gtc aca ggt gga aga aga 240
 Ser Thr Leu Thr Ser Ser Asn Arg Asn Asp Val Thr Gly Gly Arg Arg
 65 70 75 80
 gac cca aat cat tct gaa ggc tca act act tta ctg gaa ggt tat acc 288
 Asp Pro Asn His Ser Glu Gly Ser Thr Thr Leu Leu Glu Gly Tyr Thr
 85 90 95
 tct cat tac cca cac acg aag gaa agc agg acc ttc atc cca gtg acc 336
 Ser His Tyr Pro His Thr Lys Glu Ser Arg Thr Phe Ile Pro Val Thr
 100 105 110
 tca gct aag act ggg tcc ttt gga gtt act gca gtt act gtt gga gat 384
 Ser Ala Lys Thr Gly Ser Phe Gly Val Thr Ala Val Thr Val Gly Asp
 115 120 125
 tcc aac tct aat gtc aat cgt tcc tta tca gga gac caa gac aca 429
 Ser Asn Ser Asn Val Asn Arg Ser Leu Ser Gly Asp Gln Asp Thr
 130 135 140

<210> 18
 <211> 143
 <212> PRT
 <213> Homo Sapiens

<400> 18
 Asp Arg Ile Pro Ala Thr Asn Met Asp Ser Ser His Ser Thr Thr Leu
 1 5 10 15
 Gln Pro Thr Ala Asn Pro Asn Thr Gly Leu Val Glu Asp Leu Asp Arg
 20 25 30
 Thr Gly Pro Leu Ser Met Thr Thr Gln Gln Ser Asn Ser Gln Ser Phe
 35 40 45
 Ser Thr Ser His Glu Gly Leu Glu Glu Asp Lys Asp His Pro Thr Thr
 50 55 60
 Ser Thr Leu Thr Ser Ser Asn Arg Asn Asp Val Thr Gly Gly Arg Arg
 65 70 75 80
 Asp Pro Asn His Ser Glu Gly Ser Thr Thr Leu Leu Glu Gly Tyr Thr

85 90 95
 Ser His Tyr Pro His Thr Lys Glu Ser Arg Thr Phe Ile Pro Val Thr
 100 105 110
 Ser Ala Lys Thr Gly Ser Phe Gly Val Thr Ala Val Thr Val Gly Asp
 115 120 125
 Ser Asn Ser Asn Val Asn Arg Ser Leu Ser Gly Asp Gln Asp Thr
 130 135 140

<210> 19
 <211> 555
 <212> DNA
 <213> Homo Sapiens

<220>
 <221> CDS
 <222> (1)...(555)

<400> 19
 gac aga atc cct gct acc agt acg tct tca aat acc atc tca gca ggc 48
 Asp Arg Ile Pro Ala Thr Ser Thr Ser Ser Asn Thr Ile Ser Ala Gly
 1 5 10 15
 tgg gag cca aat gaa gaa agt gaa gat gaa aga gac aga cac ctc agt 96
 Trp Glu Pro Asn Glu Glu Ser Glu Asp Glu Arg Asp Arg His Leu Ser
 20 25 30
 ttt tct gga tca ggc att gat gat gat gaa gat ttt atc tcc agc acc 144
 Phe Ser Gly Ser Gly Ile Asp Asp Asp Glu Asp Phe Ile Ser Ser Thr
 35 40 45
 aat atg gac tcc agt cat agt aca acg ctt cag cct act gca aat cca 192
 Asn Met Asp Ser Ser His Ser Thr Thr Leu Gln Pro Thr Ala Asn Pro
 50 55 60
 aac aca ggt ttg gtg gaa gat ttg gac agg aca gga cct ctt tca atg 240
 Asn Thr Gly Leu Val Glu Asp Leu Asp Arg Thr Gly Pro Leu Ser Met
 65 70 75 80
 aca acg cag cag agt aat tct cag agc ttc tct aca tca cat gaa ggc 288
 Thr Thr Gln Gln Ser Asn Ser Gln Ser Phe Ser Thr Ser His Glu Gly
 85 90 95
 ttg gaa gaa gat aaa gac cat cca aca act tct act ctg aca tca agc 336
 Leu Glu Glu Asp Lys Asp His Pro Thr Thr Ser Thr Leu Thr Ser Ser
 100 105 110
 aat agg aat gat gtc aca ggt gga aga aga gac cca aat cat tct gaa 384
 Asn Arg Asn Asp Val Thr Gly Gly Arg Arg Asp Pro Asn His Ser Glu
 115 120 125
 ggc tca act act tta ctg gaa ggt tat acc tct cat tac cca cac acg 432
 Gly Ser Thr Thr Leu Leu Glu Gly Tyr Thr Ser His Tyr Pro His Thr
 130 135 140
 aag gaa agc agg acc ttc atc cca gtg acc tca gct aag act ggg tcc 480
 Lys Glu Ser Arg Thr Phe Ile Pro Val Thr Ser Ala Lys Thr Gly Ser
 145 150 155 160
 ttt gga gtt act gca gtt act gtt gga gat tcc aac tct aat gtc aat 528
 Phe Gly Val Thr Ala Val Thr Val Gly Asp Ser Asn Ser Asn Val Asn
 165 170 175
 cgt tcc tta tca gga gac caa gac aca 555
 Arg Ser Leu Ser Gly Asp Gln Asp Thr
 180 185

<210> 20
 <211> 185
 <212> PRT
 <213> Homo Sapiens

<400> 20
 Asp Arg Ile Pro Ala Thr Ser Thr Ser Ser Asn Thr Ile Ser Ala Gly
 1 5 10 15
 Trp Glu Pro Asn Glu Glu Ser Glu Asp Glu Arg Asp Arg His Leu Ser
 20 25 30
 Phe Ser Gly Ser Gly Ile Asp Asp Asp Glu Asp Phe Ile Ser Ser Thr
 35 40 45
 Asn Met Asp Ser Ser His Ser Thr Thr Leu Gln Pro Thr Ala Asn Pro
 50 55 60
 Asn Thr Gly Leu Val Glu Asp Leu Asp Arg Thr Gly Pro Leu Ser Met
 65 70 75 80
 Thr Thr Gln Gln Ser Asn Ser Gln Ser Phe Ser Thr Ser His Glu Gly
 85 90 95
 Leu Glu Glu Asp Lys Asp His Pro Thr Thr Ser Thr Leu Thr Ser Ser
 100 105 110
 Asn Arg Asn Asp Val Thr Gly Gly Arg Arg Asp Pro Asn His Ser Glu
 115 120 125
 Gly Ser Thr Thr Leu Leu Glu Gly Tyr Thr Ser His Tyr Pro His Thr
 130 135 140
 Lys Glu Ser Arg Thr Phe Ile Pro Val Thr Ser Ala Lys Thr Gly Ser
 145 150 155 160
 Phe Gly Val Thr Ala Val Thr Val Gly Asp Ser Asn Ser Asn Val Asn
 165 170 175
 Arg Ser Leu Ser Gly Asp Gln Asp Thr
 180 185

<210> 21
 <211> 561
 <212> DNA
 <213> Homo Sapiens

<220>
 <221> CDS
 <222> (1)...(561)

<400> 21
 gac aga atc cct gct acc aca gcc tca gct cat acc agc cat cca atg 48
 Asp Arg Ile Pro Ala Thr Thr Ala Ser Ala His Thr Ser His Pro Met
 1 5 10 15
 caa gga agg aca aca cca agc cca gag gac agt tcc tgg act gat ttc 96
 Gln Gly Arg Thr Thr Pro Ser Pro Glu Asp Ser Ser Trp Thr Asp Phe
 20 25 30
 ttc aac cca atc tca cac ccc atg gga cga ggt cat caa gca gga aga 144
 Phe Asn Pro Ile Ser His Pro Met Gly Arg Gly His Gln Ala Gly Arg
 35 40 45
 agg atg gat atg gac tcc agt cat agt aca acg ctt cag cct act gca 192
 Arg Met Asp Met Asp Ser Ser His Ser Thr Thr Leu Gln Pro Thr Ala
 50 55 60
 aat cca aac aca ggt ttg gtg gaa gat ttg gac agg aca gga cct ctt 240
 Asn Pro Asn Thr Gly Leu Val Glu Asp Leu Asp Arg Thr Gly Pro Leu
 65 70 75 80
 tca atg aca acg cag cag agt aat tct cag agc ttc tct aca tca cat 288
 Ser Met Thr Thr Gln Gln Ser Asn Ser Gln Ser Phe Ser Thr Ser His
 85 90 95

gaa ggc ttg gaa gaa gat aaa gac cat cca aca act tct act ctg aca 336
 Glu Gly Leu Glu Glu Asp Lys Asp His Pro Thr Thr Ser Thr Leu Thr
 100 105 110
 tca agc aat agg aat gat gtc aca ggt gga aga aga gac cca aat cat 384
 Ser Ser Asn Arg Asn Asp Val Thr Gly Gly Arg Arg Asp Pro Asn His
 115 120 125
 tct gaa ggc tca act act tta ctg gaa ggt tat acc tct cat tac cca 432
 Ser Glu Gly Ser Thr Thr Leu Leu Glu Gly Tyr Thr Ser His Tyr Pro
 130 135 140
 cac acg aag gaa agc agg acc ttc atc cca gtg acc tca gct aag act 480
 His Thr Lys Glu Ser Arg Thr Phe Ile Pro Val Thr Ser Ala Lys Thr
 145 150 155 160
 ggg tcc ttt gga gtt act gca gtt act gtt gga gat tcc aac tct aat 528
 Gly Ser Phe Gly Val Thr Ala Val Thr Val Gly Asp Ser Asn Ser Asn
 165 170 175
 gtc aat cgt tcc tta tca gga gac caa gac aca 561
 Val Asn Arg Ser Leu Ser Gly Asp Gln Asp Thr
 180 185

<210> 22
 <211> 187
 <212> PRT
 <213> Homo Sapiens

<400> 22
 Asp Arg Ile Pro Ala Thr Thr Ala Ser Ala His Thr Ser His Pro Met
 1 5 10 15
 Gln Gly Arg Thr Pro Ser Pro Glu Asp Ser Ser Trp Thr Asp Phe
 20 25 30
 Phe Asn Pro Ile Ser His Pro Met Gly Arg Gly His Gln Ala Gly Arg
 35 40 45
 Arg Met Asp Met Asp Ser Ser His Ser Thr Thr Leu Gln Pro Thr Ala
 50 55 60
 Asn Pro Asn Thr Gly Leu Val Glu Asp Leu Asp Arg Thr Gly Pro Leu
 65 70 75 80
 Ser Met Thr Thr Gln Gln Ser Asn Ser Gln Ser Phe Ser Thr Ser His
 85 90 95
 Glu Gly Leu Glu Glu Asp Lys Asp His Pro Thr Thr Ser Thr Leu Thr
 100 105 110
 Ser Ser Asn Arg Asn Asp Val Thr Gly Gly Arg Arg Asp Pro Asn His
 115 120 125
 Ser Glu Gly Ser Thr Thr Leu Leu Glu Gly Tyr Thr Ser His Tyr Pro
 130 135 140
 His Thr Lys Glu Ser Arg Thr Phe Ile Pro Val Thr Ser Ala Lys Thr
 145 150 155 160
 Gly Ser Phe Gly Val Thr Ala Val Thr Val Gly Asp Ser Asn Ser Asn
 165 170 175
 Val Asn Arg Ser Leu Ser Gly Asp Gln Asp Thr
 180 185

<210> 23
 <211> 193
 <212> PRT
 <213> Rattus Norvegicus

<400> 23
 Gln Leu Arg Asn Phe Leu Lys Cys Ser Glu Asp Asn Pro Leu Phe Ala
 1 5 10 15
 Gly Ile Asp Cys Glu Val Phe Glu Ser Arg Phe Pro Thr Thr Met Ala
 20 25 30

Leu Ser Val³⁵ Leu Val Thr Ile⁴⁰ Glu Met Cys Asn Ala⁴⁵ Leu Asn Ser Val
 Ser Glu Asn Gln Ser Leu⁵⁵ Leu Arg Met Pro Pro Trp⁶⁰ Leu Asn Pro Trp
 Leu Leu Gly Ala Val⁷⁰ Val Met Ser Met Ala⁷⁵ Leu His Phe Leu Ile⁸⁰ Leu
 Leu Val Pro Pro⁸⁵ Leu Pro Leu Ile Phe⁹⁰ Gln Val Thr Pro Leu Ser Gly⁹⁵
 Arg Gln Trp Gly¹⁰⁰ Val Val Leu Gln Met¹⁰⁵ Ser Leu Pro Val Ile¹¹⁰ Leu Leu
 Asp Glu Ala¹¹⁵ Leu Lys Tyr Leu Ser Arg His His Val¹²⁵ Asp Gly Val Leu
 Glu Thr Phe Met Gln Ala¹³⁵ Trp Cys Lys Gln Pro¹⁴⁰ Leu Pro Gly Pro His
 Thr¹³⁰ Thr Arg Gly Trp Leu¹⁵⁰ Pro Gly Cys His Phe¹⁵⁵ Asn Gly Trp Glu Gln¹⁶⁰
 Thr¹⁴⁵ Glu Glu Phe Val¹⁶⁵ Phe Ile Gln Glu Arg¹⁷⁰ Trp Thr Val Ser Gly¹⁷⁵ Leu
 Gly Pro Glu Lys¹⁸⁰ Lys Ala Arg Glu Arg¹⁸⁵ Leu Gly Leu Val Ser¹⁹⁰ Ala Ala
 Ser

<210> 24
 <211> 161
 <212> PRT
 <213> Homo Sapiens

<400> 24
 Glu Ser Arg Phe⁵ Thr Thr Met Ala¹⁰ Leu Ser Val Leu Val Thr Ile¹⁵
 Glu¹ Met Cys Asn Ala Leu Asn Ser Val²⁵ Ser Glu Asn Gln Ser³⁰ Leu Leu
 Arg Met Pro³⁵ Pro Trp Met Asn Pro⁴⁰ Trp Leu Leu Val Ala⁴⁵ Val Ala Met
 Ser Met Ala Leu His Phe⁵⁵ Leu Ile Leu Leu Val Pro⁶⁰ Pro Leu Pro Leu
 Ile⁵⁰ Phe Gln Val Thr⁷⁰ Pro Leu Ser Gly Arg Gln Trp Val Val Val Leu⁸⁰
 Gln Ile Ser Leu⁸⁵ Val Ile Leu Leu Asp⁹⁰ Glu Ala Leu Lys Tyr⁹⁵ Leu
 Ser Arg Asn His¹⁰⁰ Met His Ala Cys Leu¹⁰⁵ Tyr Pro Gly Leu Leu Arg Thr¹¹⁰
 Val Ser Gln Ala Trp Ser Arg Gln¹²⁰ Pro Leu Thr Thr Ser Trp Thr Pro¹²⁵
 Asp His Thr Gly Leu Ala Ser¹³⁵ Leu Gly Gln Gly His Ser Ile Val Ser¹⁴⁰
 Leu¹³⁰ Ser Glu Leu Leu Arg¹⁵⁰ Glu Gly Gly Ser Arg¹⁵⁵ Glu Glu Met Ser Gln¹⁶⁰
 Lys¹⁴⁵

<210> 25
 <211> 153
 <212> PRT
 <213> Homo sapiens

<400> 25
 Glu Ser Arg Phe⁵ Thr Thr Met Ala¹⁰ Leu Ser Val Leu Val Thr Ile¹⁵
 Glu¹ Met Cys Asn Ala Leu Asn Ser Val²⁵ Ser Glu Asn Gln Ser³⁰ Leu Leu
 Arg Met Pro³⁵ Pro Trp Met Asn Pro⁴⁰ Trp Leu Leu Val Ala⁴⁵ Val Ala Met
 Ser Met Ala Leu His Phe Leu Ile Leu Leu Val Pro Leu Pro Leu

```

      50      55      60
Ile Phe Gln Val Thr Pro Leu Ser Gly Arg Gln Trp Val Val Val Leu
65      70      75      80
Gln Ile Ser Leu Pro Val Ile Leu Leu Asp Glu Ala Leu Lys Tyr Leu
      85      90      95
Ser Arg Asn His Met His Ala Cys Leu Tyr Pro Gly Leu Leu Arg Thr
      100      105      110
Val Ser Gln Ala Trp Ser Arg Gln Pro Leu Thr Thr Ser Trp Thr Pro
      115      120      125
Asp His Thr Gly Ala Arg Asp Thr Ala Ser Ser Arg Cys Gln Ser Cys
      130      135      140
Ser Glu Arg Glu Glu Ala Gly Lys Lys
145      150

```

<210> 26
 <211> 17
 <212> PRT
 <213> Homo sapiens

```

<400> 26
Leu Ile Leu Val Gly Glu Pro Ser Ile Ser Thr Pro Asp Gly Thr Ile
1      5      10      15
Lys

```

<210> 27
 <211> 101
 <212> PRT
 <213> Homo sapiens

```

<400> 27
Met Glu Asp Asp Glu Leu Thr Asp Ser Lys Leu Pro Ser His Ala Thr
1      5      10      15
His Ser Leu Phe Thr Cys Pro Glu Asn Glu Glu Met Val Leu Ser Asn
      20      25      30
Ser Arg Ile Gly Lys Arg Arg Gly Glu Pro Leu Ile Leu Val Gly Glu
      35      40      45
Pro Ser Ile Ser Thr Pro Asp Gly Thr Ile Lys Asp Arg Arg Leu Phe
      50      55      60
Met His His Val Ser Leu Glu Pro Ile Thr Cys Val Pro Phe Arg Thr
65      70      75      80
Thr Lys Glu Arg Gln Glu Ile Gln Asn Pro Asn Phe Thr Ala Pro Gly
      85      90      95
Gln Glu Phe Leu Ser
      100

```

<210> 28
 <211> 149
 <212> PRT
 <213> Homo sapiens

```

<400> 28
Ala Glu Lys Gln Ala Lys Ala Arg Glu Ser Gly Ser Ser Thr Ala Glu
1      5      10      15
Glu Gly Asp Phe Ser Lys Gly Pro Ile Arg Cys Asp Arg Cys Gly Tyr
      20      25      30
Asn Thr Asn Arg Tyr Asp His Tyr Thr Ala His Leu Lys His His Thr
      35      40      45
Arg Ala Gly Asp Asn Glu Arg Val Tyr Lys Cys Ile Ile Cys Thr Tyr
      50      55      60
Thr Thr Val Ser Glu Tyr His Trp Arg Lys His Leu Arg Asn His Phe
65      70      75      80
Pro Arg Lys Val Tyr Thr Cys Gly Lys Cys Asn Tyr Phe Ser Asp Arg
      85      90      95

```

Lys Asn Asn Tyr Val Gln His Val Arg Thr His Thr Gly Glu Arg Pro
 100 105 110
 Tyr Lys Cys Glu Leu Cys Pro Tyr Ser Ser Ser Gln Lys Thr His Leu
 115 120 125
 Thr Arg His Met Arg Thr His Ser Val Gly Tyr Gly Tyr His Leu Val
 130 135 140
 Ile Phe Thr Arg Val
 145

<210> 29
 <211> 13
 <212> PRT
 <213> Homo sapiens

<400> 29
 Val Gly Tyr Gly Tyr His Leu Val Ile Phe Thr Arg Val
 1 5 10

<210> 30
 <211> 875
 <212> DNA
 <213> Homo sapiens

<400> 30
 cgcttgccg agctgtacgt gaagccgggc aacaaggaac gcggttgaa cgacccgccg 60
 cagttctcat acgggctgca gaccagggc ggcggacca ggcgtcgct gcttaccag 120
 agggtagccg caccacagga tggatcccc agagtcccc catcagagac ttctcctggg 180
 cctcccccaa tggggcctcc acctccttca agtaaggctc ccagggtccc acctgtgggg 240
 agtggctcctg cctctggcgt ggagcccaca agtttcccag tcgagtctga ggctgtgatg 300
 gaggatgtgc tgagaccttt ggaacaggca ttggaagact gccgtggcca cacaaggaag 360
 caggtatgtg atgacatcag ccgacgcctg gcactgctgc aggaacagt ggctggagg 420
 aagttgtcaa tacctgtaaa gaagagaatg gctctactgg tgcaagagct ttcaagccac 480
 cggtgggacg cagcagatga catccaccgc tccctcatgg ttgacctgt gactgaggtc 540
 agtcagtggg tggtaggagt taaaagatta attgcagaaa agaggagtct gttttcagag 600
 gaggcagcca atgaagagaa atctgcagcc acagctgaga agaaccatac cataccaggc 660
 ttccagcagg ctctataatc ctcggttccc cagactcacc ggacaccatc tcctatgcct 720
 tggagacctt ctgtcacttg gctcccttct taccaccacc aagactgtcc cactgggcct 780
 gaccaccta tgagggaaga agtcccacct gggccagagg gagttcatgt gttactcata 840
 acatgcattt caataaaaac atctctgcgg tgggtg 875

<210> 31
 <211> 672
 <212> DNA
 <213> Homo sapiens

<400> 31
 cgcttgccg agctgtacgt gaagccgggc aacaaggaac gcggttgaa cgacccgccg 60
 cagttctcat acgggctgca gaccagggc ggcggacca ggcgtcgct gcttaccag 120
 agggtagccg caccacagga tggatcccc agagaagcag gtatgtgatg acatcagccg 180
 acgcctggca ctgctgcagg aacagtgggc tggaggaaaag ttgtcaatac ctgtaaaaga 240
 gagaatggct ctactggtgc aagagctttc aagccaccgg tgggacgcag cagatgacat 300
 ccaccgtccc ctcatggttg accatgtgac tgagggtcagt cagtggatgg taggagttaa 360
 aagattaatt gcagaaaaga ggagtctggt ttcagaggag gcagccaatg aagagaaatc 420
 tgcagccaca gctgagaaga accataccat accaggcttc cagcaggctt cataatcctc 480
 gggtccccag actcaccgga caccatctcc tatgccttgg agaccttctg tcacttggct 540
 cccttcttac caccaccaag actgtcccac tgggcctgac ccacctatga gggaagaagt 600
 cccacctggg ccagagggag ttcattgtgt actcataaca tgcatttcaa taaaaacatc 660
 tctgcgggtg tg 672

<210> 32
 <211> 598
 <212> PRT
 <213> Homo sapiens

<400> 32

Met Ser Asp Asn Gln Ser Trp Asn Ser Ser Gly Ser Glu Glu Asp Pro
 1 5 10 15
 Glu Thr Glu Ser Gly Pro Pro Val Glu Arg Cys Gly Val Leu Ser Lys
 20 25 30
 Trp Thr Asn Tyr Ile His Gly Trp Gln Asp Arg Trp Val Val Leu Lys
 35 40 45
 Asn Asn Ala Leu Ser Tyr Tyr Lys Ser Glu Asp Glu Thr Glu Tyr Gly
 50 55 60
 Cys Arg Gly Ser Ile Cys Leu Ser Lys Ala Val Ile Thr Pro His Asp
 65 70 75 80
 Phe Asp Glu Cys Arg Phe Asp Ile Ser Val Asn Asp Ser Val Trp Tyr
 85 90 95
 Leu Arg Ala Gln Asp Pro Asp His Arg Gln Gln Trp Ile Asp Ala Ile
 100 105 110
 Glu Gln His Lys Thr Glu Ser Gly Tyr Gly Ser Glu Ser Ser Leu Arg
 115 120 125
 Arg His Gly Ser Met Val Ser Leu Val Ser Gly Ala Ser Gly Tyr Ser
 130 135 140
 Ala Thr Ser Thr Ser Ser Phe Lys Lys Gly His Ser Leu Arg Glu Lys
 145 150 155 160
 Leu Ala Glu Met Glu Thr Phe Arg Asp Ile Leu Cys Arg Gln Val Asp
 165 170 175
 Thr Leu Gln Lys Tyr Phe Asp Ala Cys Ala Asp Ala Val Ser Lys Asp
 180 185 190
 Glu Leu Gln Arg Asp Lys Val Val Glu Asp Asp Glu Asp Asp Phe Pro
 195 200 205
 Thr Thr Arg Ser Asp Gly Asp Phe Leu His Ser Thr Asn Gly Asn Lys
 210 215 220
 Glu Lys Leu Phe Pro His Val Thr Pro Lys Gly Ile Asn Gly Ile Asp
 225 230 235 240
 Phe Lys Gly Glu Ala Ile Thr Phe Lys Ala Thr Thr Ala Gly Ile Leu
 245 250 255
 Ala Thr Leu Ser His Cys Ile Glu Leu Met Val Lys Arg Glu Asp Ser
 260 265 270
 Trp Gln Lys Arg Leu Asp Lys Glu Thr Glu Lys Lys Arg Arg Thr Glu
 275 280 285
 Glu Ala Tyr Lys Asn Ala Met Thr Glu Leu Lys Lys Lys Ser His Phe
 290 295 300
 Gly Gly Pro Asp Tyr Glu Glu Gly Pro Asn Ser Leu Ile Asn Glu Glu
 305 310 315 320
 Glu Phe Phe Asp Ala Val Glu Ala Ala Leu Asp Arg Gln Asp Lys Ile
 325 330 335
 Glu Glu Gln Ser Gln Ser Glu Lys Val Arg Leu His Trp Pro Thr Ser
 340 345 350
 Leu Pro Ser Gly Asp Ala Phe Ser Ser Val Gly Thr His Arg Phe Val
 355 360 365
 Gln Lys Val Glu Glu Met Val Gln Asn His Met Thr Tyr Ser Leu Gln
 370 375 380
 Asp Val Gly Gly Asp Ala Asn Trp Gln Leu Val Val Glu Glu Gly Glu
 385 390 395 400
 Met Lys Val Tyr Arg Arg Glu Val Glu Glu Asn Gly Ile Val Leu Asp
 405 410 415
 Pro Leu Lys Ala Thr His Ala Val Lys Gly Val Thr Gly His Glu Val
 420 425 430
 Cys Asn Tyr Phe Trp Asn Val Asp Val Arg Asn Asp Trp Glu Thr Thr
 435 440 445
 Ile Glu Asn Phe His Val Val Glu Thr Leu Ala Asp Asn Ala Ile Ile
 450 455 460
 Ile Tyr Gln Thr His Lys Arg Val Trp Pro Ala Ser Gln Arg Asp Val
 465 470 475 480
 Leu Tyr Leu Ser Val Ile Arg Lys Ile Pro Ala Leu Thr Glu Asn Asp
 485 490 495
 Pro Glu Thr Trp Ile Val Cys Asn Phe Ser Val Asp His Asp Ser Ala
 500 505 510
 Pro Leu Asn Asn Arg Cys Val Arg Ala Lys Ile Asn Val Ala Met Ile
 515 520 525
 Cys Gln Thr Leu Val Ser Pro Pro Glu Gly Asn Gln Glu Ile Ser Arg

530	Asp	Asn	Ile	Leu	Cys	Lys	535	Ile	Thr	Tyr	Val	Ala	540	Asn	Val	Asn	Pro	Gly
545	Gly	Trp	Ala	Pro	Ala	Ser	550	Val	Leu	Arg	Ala	Val	555	Ala	Lys	Arg	Glu	Tyr
	Pro	Lys	Phe	Leu	Lys	Arg	565	Phe	Thr	Ser	570	Val	Gln	Glu	Lys	Thr	Ala	
	Gly	Lys	Pro	580	Ile	Leu	Phe			585					590			
			595															

<210> 33
 <211> 572
 <212> PRT
 <213> Homo sapiens

<400> 33

Met	Ser	Asp	Asn	Gln	Ser	Trp	Asn	Ser	Ser	Gly	Ser	Glu	Glu	Asp	Pro
1				5					10					15	
Glu	Thr	Glu	Ser	Gly	Pro	Pro	Val	Glu	Arg	Cys	Gly	Val	Leu	Ser	Lys
			20					25					30		
Trp	Thr	Asn	Tyr	Ile	His	Gly	Trp	Gln	Asp	Arg	Trp	Val	Val	Leu	Lys
		35					40					45			
Asn	Asn	Ala	Leu	Ser	Tyr	Tyr	Lys	Ser	Glu	Asp	Glu	Thr	Glu	Tyr	Gly
	50					55					60				
Cys	Arg	Gly	Ser	Ile	Cys	Leu	Ser	Lys	Ala	Val	Ile	Thr	Pro	His	Asp
65					70					75					80
Phe	Asp	Glu	Cys	Arg	Phe	Asp	Ile	Ser	Val	Asn	Asp	Ser	Val	Trp	Tyr
				85					90					95	
Leu	Arg	Ala	Gln	Asp	Pro	Asp	His	Arg	Gln	Gln	Trp	Ile	Asp	Ala	Ile
			100					105					110		
Glu	Gln	His	Lys	Thr	Glu	Ser	Gly	Tyr	Gly	Ser	Glu	Ser	Ser	Leu	Arg
		115					120					125			
Arg	His	Gly	Ser	Met	Val	Ser	Leu	Val	Ser	Gly	Ala	Ser	Gly	Tyr	Ser
	130					135					140				
Ala	Thr	Ser	Thr	Ser	Ser	Phe	Lys	Lys	Gly	His	Ser	Leu	Arg	Glu	Lys
145					150					155					160
Leu	Ala	Glu	Met	Glu	Thr	Phe	Arg	Asp	Ile	Leu	Cys	Arg	Gln	Val	Asp
				165					170					175	
Thr	Leu	Gln	Lys	Tyr	Phe	Asp	Ala	Cys	Ala	Asp	Ala	Val	Ser	Lys	Asp
			180					185					190		
Glu	Leu	Gln	Arg	Asp	Lys	Val	Val	Glu	Asp	Asp	Glu	Asp	Asp	Phe	Pro
		195					200					205			
Thr	Thr	Arg	Ser	Asp	Gly	Asp	Phe	Leu	His	Ser	Thr	Asn	Gly	Asn	Lys
	210					215						220			
Glu	Lys	Leu	Phe	Pro	His	Val	Thr	Pro	Lys	Gly	Ile	Asn	Gly	Ile	Asp
225					230					235					240
Phe	Lys	Gly	Glu	Ala	Ile	Thr	Phe	Lys	Ala	Thr	Thr	Ala	Gly	Ile	Leu
				245					250					255	
Ala	Thr	Leu	Ser	His	Cys	Ile	Glu	Leu	Met	Val	Lys	Arg	Glu	Asp	Ser
			260					265					270		
Trp	Gln	Lys	Arg	Leu	Asp	Lys	Glu	Thr	Glu	Lys	Lys	Arg	Arg	Thr	Glu
		275					280					285			
Glu	Ala	Tyr	Lys	Asn	Ala	Met	Thr	Glu	Leu	Lys	Lys	Lys	Ser	His	Phe
	290					295					300				
Gly	Gly	Pro	Asp	Tyr	Glu	Glu	Gly	Pro	Asn	Ser	Leu	Ile	Asn	Glu	Glu
305					310					315					320
Glu	Phe	Phe	Asp	Ala	Val	Glu	Ala	Ala	Leu	Asp	Arg	Gln	Asp	Lys	Ile
				325					330					335	
Glu	Glu	Gln	Ser	Gln	Ser	Glu	Lys	Val	Arg	Leu	His	Trp	Pro	Thr	Ser
			340					345					350		
Leu	Pro	Ser	Gly	Asp	Ala	Phe	Ser	Ser	Val	Gly	Thr	His	Arg	Phe	Val
		355					360					365			
Gln	Lys	Glu	Glu	Gly	Glu	Met	Lys	Val	Tyr	Arg	Arg	Glu	Val	Glu	Glu
	370					375					380				
Asn	Gly	Ile	Val	Leu	Asp	Pro	Leu	Lys	Ala	Thr	His	Ala	Val	Lys	Gly
385					390					395					400

Val Thr Gly His Glu Val Cys Asn Tyr Phe Trp Asn Val Asp Val Arg
 405 410 415
 Asn Asp Trp Glu Thr Thr Ile Glu Asn Phe His Val Val Glu Thr Leu
 420 425 430
 Ala Asp Asn Ala Ile Ile Ile Tyr Gln Thr His Lys Arg Val Trp Pro
 435 440 445
 Ala Ser Gln Arg Asp Val Leu Tyr Leu Ser Val Ile Arg Lys Ile Pro
 450 455 460
 Ala Leu Thr Glu Asn Asp Pro Glu Thr Trp Ile Val Cys Asn Phe Ser
 465 470 475 480
 Val Asp His Asp Ser Ala Pro Leu Asn Asn Arg Cys Val Arg Ala Lys
 485 490 495
 Ile Asn Val Ala Met Ile Cys Gln Thr Leu Val Ser Pro Pro Glu Gly
 500 505 510
 Asn Gln Glu Ile Ser Arg Asp Asn Ile Leu Cys Lys Ile Thr Tyr Val
 515 520 525
 Ala Asn Val Asn Pro Gly Gly Trp Ala Pro Ala Ser Val Leu Arg Ala
 530 535 540
 Val Ala Lys Arg Glu Tyr Pro Lys Phe Leu Lys Arg Phe Thr Ser Tyr
 545 550 555 560
 Val Gln Glu Lys Thr Ala Gly Lys Pro Ile Leu Phe
 565 570

<210> 34
 <211> 89
 <212> PRT
 <213> Homo sapiens

<400> 34
 Ser Gln Ser Glu Lys Val Arg Leu His Trp Pro Thr Ser Leu Pro Ser
 1 5 10 15
 Gly Asp Ala Phe Ser Ser Val Gly Thr His Arg Phe Val Gln Lys Pro
 20 25 30
 Tyr Ser Arg Ser Ser Ser Met Ser Ile Asp Leu Val Ser Ala Ser
 35 40 45
 Asp Asp Val His Arg Phe Ser Ser Gln Val Glu Glu Met Val Gln Asn
 50 55 60
 His Met Thr Tyr Ser Leu Gln Asp Val Gly Gly Asp Ala Asn Trp Gln
 65 70 75 80
 Leu Val Val Glu Glu Gly Glu Met Lys
 85

<210> 35
 <211> 63
 <212> PRT
 <213> Homo sapiens

<400> 35
 Gly Leu Pro Leu Ala Glu Ser Leu Lys Arg Leu Met Ser Leu Ser Pro
 1 5 10 15
 Gly Arg Pro Pro Leu Leu Leu Trp Asp Ala His Val Ala Asp Arg Asp
 20 25 30
 His Leu Cys Gly Gly Ser Ala His Arg Leu Thr His His Leu Glu Glu
 35 40 45
 Asp Gly Leu Arg Pro Pro Ala Ala Leu Asp Cys Val Phe Pro Pro
 50 55 60

<210> 36
 <211> 13
 <212> PRT
 <213> Homo sapiens

<400> 36
 His Leu Asp Ala Gly Thr Val Glu Pro Lys Arg Glu Lys
 Page 19

<210> 37
 <211> 57
 <212> PRT
 <213> Homo sapiens

<400> 37
 Ile Glu Thr Arg Ser Lys Asn Phe Ser Ala Cys Leu Glu Leu Gly Glu
 1 5 10 15
 Ser Leu Leu Gln Arg Gln His Gln Ala Ser Glu Glu Ile Arg Glu Lys
 20 25 30
 Leu Gln Gln Val Met Ser Arg Arg Lys Glu Met Asn Glu Lys Trp Glu
 35 40 45
 Ala Arg Trp Glu Arg Leu Arg Met Ser
 50 55

<210> 38
 <211> 10
 <212> PRT
 <213> Homo sapiens

<400> 38
 Gly Pro Pro Gly Ala Pro Gly Ala Pro Gly
 1 5 10

<210> 39
 <211> 14
 <212> PRT
 <213> Homo sapiens

<400> 39
 Thr Val Leu Leu Arg Leu Gly Ile Thr Trp Gly Lys Val Val
 1 5 10

<210> 40
 <211> 16
 <212> DNA
 <213> Homo sapiens - Alt1

<400> 40
 tctgtacctg atactc

16

<210> 41
 <211> 16
 <212> DNA
 <213> Homo sapiens - Alt2

<400> 41
 ctacagactg atactc

16

<210> 42
 <211> 11
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Protein Transduction Domain

<400> 42
 Tyr Gly Arg Lys Lys Arg Arg Gln Arg Arg
 1 5 10

<210> 43
<211> 21
<212> DNA
<213> Artificial Sequence

<220>
<223> Primer

<400> 43
ccgaaaccat gaactttctg c

21

<210> 44
<211> 21
<212> DNA
<213> Artificial Sequence

<220>
<223> Primer

<400> 44
cttggcgatt tagcagcaga t

21

<210> 45
<211> 21
<212> DNA
<213> Artificial Sequence

<220>
<223> Primer

<400> 45
accctggcctt tactgctgta c

21

<210> 46
<211> 21
<212> DNA
<213> Artificial Sequence

<220>
<223> Primer

<400> 46
aaatggcgaa tccagtccca c

21

<210> 47
<211> 21
<212> DNA
<213> Artificial Sequence

<220>
<223> Primer

<400> 47
ttactgctgt acctccacca t

21

<210> 48
<211> 21
<212> DNA
<213> Artificial Sequence

<220>
<223> Primer

<400> 48
gaaggatctc ctcttccttc a

21

<210> 49
<211> 21
<212> DNA
<213> Artificial Sequence

<220>
<223> Primer

<400> 49
ctgcttggtg cactgctgca g

21

<210> 50
<211> 21
<212> DNA
<213> Artificial Sequence

<220>
<223> Primer

<400> 50
tctggaaagc agcttgctcac t

21

<210> 51
<211> 21
<212> DNA
<213> Artificial Sequence

<220>
<223> Primer

<400> 51
gcccctgtgt cccagtttga t

21

<210> 52
<211> 21
<212> DNA
<213> Artificial Sequence

<220>
<223> Primer

<400> 52
tacaggtgac tgggttgagc t

21

<210> 53
<211> 21
<212> DNA
<213> Artificial Sequence

<220>
<223> Primer

<400> 53
agccaccaga agaaagtggt g

21

<210> 54
<211> 21
<212> DNA
<213> Artificial Sequence

<220>
<223> Primer

<400> 54
tgtctgggtt gagctctaag c

21

<210> 55

<211> 21
<212> DNA
<213> Artificial Sequence

<220>
<223> Primer

<400> 55
aacatgcact tgctgtgctt c 21

<210> 56
<211> 21
<212> DNA
<213> Artificial Sequence

<220>
<223> Primer

<400> 56
ctctcccgca gtaatccaca t 21

<210> 57
<211> 21
<212> DNA
<213> Artificial Sequence

<220>
<223> Primer

<400> 57
gaggtcaagg cttttgaagg c 21

<210> 58
<211> 21
<212> DNA
<213> Artificial Sequence

<220>
<223> Primer

<400> 58
cttgggcctc tgttaccatg t 21

<210> 59
<211> 21
<212> DNA
<213> Artificial Sequence

<220>
<223> Primer

<400> 59
gcttttgaag gcaaagacct g 21

<210> 60
<211> 21
<212> DNA
<213> Artificial Sequence

<220>
<223> Primer

<400> 60
ttaccatgtg gtccacaga g 21

<210> 61
<211> 21

<212> DNA
<213> Artificial Sequence

<220>
<223> Primer

<400> 61
ggagaatgcc ttttgcaaca c 21

<210> 62
<211> 21
<212> DNA
<213> Artificial Sequence

<220>
<223> Primer

<400> 62
gccattgcat ggaaatgtgg c 21

<210> 63
<211> 21
<212> DNA
<213> Artificial Sequence

<220>
<223> Primer

<400> 63
caactgctta gtcacggta g 21

<210> 64
<211> 21
<212> DNA
<213> Artificial Sequence

<220>
<223> Primer

<400> 64
acttgacaaa gcagtgcgt g 21

<210> 65
<211> 21
<212> DNA
<213> Artificial Sequence

<220>
<223> Primer

<400> 65
atgtatggag aatggggaat g 21

<210> 66
<211> 21
<212> DNA
<213> Artificial Sequence

<220>
<223> Primer

<400> 66
gttgaatcaa gggttctcct g 21

<210> 67
<211> 21
<212> DNA

<213> Artificial Sequence

<220>

<223> Primer

<400> 67

tctcctctgg tatcagcgtc t

21

<210> 68

<211> 21

<212> DNA

<213> Artificial Sequence

<220>

<223> Primer

<400> 68

gcactgaatt cctgagtgtc t

21

<210> 69

<211> 21

<212> DNA

<213> Artificial Sequence

<220>

<223> Primer

<400> 69

tggtgattgt gccttgaagg a

21

<210> 70

<211> 21

<212> DNA

<213> Artificial Sequence

<220>

<223> Primer

<400> 70

tccatgcccc ttatcatgga g

21

<210> 71

<211> 21

<212> DNA

<213> Artificial Sequence

<220>

<223> Primer

<400> 71

tgaaggacct tggctctgga t

21

<210> 72

<211> 21

<212> DNA

<213> Artificial Sequence

<220>

<223> Primer

<400> 72

aatagaggggt aggtaccagc a

21

<210> 73

<211> 12

<212> PRT

<213> Artificial Sequence

<220>

<223> VEGF alt splice #1

<400> 73

Arg Thr Lys Pro Glu Lys Cys Asp Lys Pro Arg Arg
1 5 10

<210> 74

<211> 36

<212> DNA

<213> Artificial Sequence

<220>

<223> VEGF alt splice #1 nucleotide

<400> 74

agaacaaagc cagaaaaatg tgacaagcca aggcgg

36

<210> 75

<211> 13

<212> PRT

<213> Artificial Sequence

<220>

<223> VEGFB alt splice #1 peptide

<400> 75

Val Val Lys Gln Leu Val Gln Thr Pro Pro Leu Pro Pro
1 5 10

<210> 76

<211> 39

<212> DNA

<213> Artificial Sequence

<220>

<223> VEGFB alt splice #1 nucleotide

<400> 76

gtggtcaaac aactagtgca gacgccgccg cttcctcca

39

<210> 77

<211> 12

<212> PRT

<213> Artificial Sequence

<220>

<223> VEGFD alt splice #1 peptide

<400> 77

His Gly Pro Val Lys Met Ser Ser Phe Gln Glu Thr
1 5 10

<210> 78

<211> 36

<212> DNA

<213> Artificial Sequence

<220>

<223> VEGFD alt splice #1 nucleotide

<400> 78

catggaccag tgaagatgtc ctcatccaa gaaact

36

<210> 79
<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> VEGFD alt splice #2 peptide

<400> 79
Leu Glu Arg Ser Glu Ser Cys Glu Asp Arg Cys Pro
1 5 10

<210> 80
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> VEGFD alt splice #2 nucleotide

<400> 80
ttggaacgat ctgaaagctg tgaggacaga tgtcct 36

<210> 81
<211> 13
<212> PRT
<213> Artificial Sequence

<220>
<223> VEGF alt splice #1 peptide + C

<400> 81
Arg Thr Lys Pro Glu Lys Cys Asp Lys Pro Arg Arg Cys
1 5 10

<210> 82
<211> 14
<212> PRT
<213> Artificial Sequence

<220>
<223> EGFRVIII peptide

<400> 82
Leu Glu Glu Lys Lys Gly Asn Tyr Val Val Thr Asp His Cys
1 5 10

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
5 December 2002 (05.12.2002)

PCT

(10) International Publication Number
WO 02/097044 A3

(51) International Patent Classification⁷: **A61K 38/00**,
35/34, 38/24, C12P 21/06

(21) International Application Number: PCT/US02/16707

(22) International Filing Date: 28 May 2002 (28.05.2002)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/293,791 25 May 2001 (25.05.2001) US

(71) Applicant (*for all designated States except US*):
THOMAS JEFFERSON UNIVERSITY [US/US];
1020 Walnut Street, Philadelphia, PA 19107-5587 (US).

(72) Inventor; and

(75) Inventor/Applicant (*for US only*): **WONG, Albert, J.**
[US/US]; Apartment 3513, 1500 Locust Street, Philadel-
phia, PA 19103 (US).

(74) Agent: **MONACO, Daniel, A.**; Drinker, Biddle & Reath,
LLP., One Logan Square, 18th and Cherry Streets, Philadel-
phia, PA 19103-6996 (US).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,
MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG,
SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ,
VN, YU, ZA, ZM, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR,
GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent
(BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR,
NE, SN, TD, TG).

Published:

— with international search report

(88) Date of publication of the international search report:
28 August 2003

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: ALTERNATIVE SPLICE FORMS OF PROTEINS AS BASIS FOR MULTIPLE THERAPEUTIC MODALITIES

(57) Abstract: Peptides or antibodies derived from alternative splice forms of proteins associated with a disease or physiologic condition are used as therapeutic or prophylactic agents. Peptides or antibodies derived from alternative splice forms of the vascular endothelial growth factor (VEGF) family of proteins are particularly useful in preventing or delaying the onset of tumors and inducing tumor regression.

WO 02/097044 A3

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US02/16707

A. CLASSIFICATION OF SUBJECT MATTER

IPC(7) : A61K 38/00, 35/34, 38/24; C12P 21/06

US CL : 514/12; 424/569; 435/252.3, 320.1, 69.1; 530/350,399

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHEDMinimum documentation searched (classification system followed by classification symbols)
U.S. : 514/12; 424/569; 435/252.3, 320.1, 69.1; 530/350,399

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)
SEQ SEARCH, WEST, STN CAS: MEDLINE CAPLUS EMBASE BIOSIS**C. DOCUMENTS CONSIDERED TO BE RELEVANT**

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X — Y	US 5,994,300 A (BAYNE et al) 30 November 1999 (30.11.1999), entire patent.	1, 5-7, 17, 25-26, 30-32, 50-63, 66-72 2-4, 8-16, 18-24, 27-29, 33-49, 64-65 and 73-76
X — Y	US 5,726,152 A (BAYNE et al) 10 March 1998 (10.03.1998), entire patent.	1, 5-7, 17, 25-26, 30-32, 50-63, 66-72 2-4, 8-16, 18-24, 27-29, 33-49, 64-65, 73-76

☐ Further documents are listed in the continuation of Box C.☐ See patent family annex.

Special categories of cited documents:	
* "A" document defining the general state of the art which is not considered to be of particular relevance	"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
"E" earlier application or patent published on or after the international filing date	"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
"O" document referring to an oral disclosure, use, exhibition or other means	"&" document member of the same patent family
"P" document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

04 May 2003 (04.05.2003)

Date of mailing of the international search report

15 MAY 2003

Name and mailing address of the ISA/US

Mail Stop PCT, Attn: ISA/US
Commissioner for Patents
P.O. Box 1450
Alexandria, Virginia 22313-1450

Facsimile No. (703)305-3230

Authorized officer

Amy DeCloux

Telephone No. 703 308-0196

Form PCT/ISA/210 (second sheet) (July 1998)